

Connectivity of the Coset Poset

Daniel A. Ramras

Department of Mathematics

Cornell University

May, 2002

(Revised September, 2002)

Contents

1	Introduction	3
2	Background Material	6
2.1	Simplicial Complexes	6
2.2	Homotopy Theory for Posets	11
3	Connectivity of the Coset Poset	24
3.1	Conditions for Simple Connectivity	24
3.2	Mayer-Vietoris Sequences and $H_*(\mathcal{C}(G))$	31
3.3	Applications to Group Theory	37
3.4	Group Extensions	38
4	Examples	41
4.1	The Quaternion Group	41
4.2	The Coset Poset of A_5	43
4.3	The Coset Poset of $PSL_2(\mathbb{F}_7)$	48
4.3.1	The Subgroups of $PSL_2(\mathbb{F}_p)$	51
4.3.2	The Möbius Function of $PSL_2(\mathbb{F}_7)$	65
4.3.3	Simple Connectivity of $\mathcal{C}(PSL_2(\mathbb{F}_7))$	71
5	Conjectures and Directions for Further Research	77

Acknowledgements:

I would like to thank Professor Kenneth S. Brown for his guidance throughout the course of this project. Professor Brown served as my mentor for the two years I spent on this research, and his assistance was invaluable. In particular, Professor Brown suggested studying the coset poset and introduced me to most of the techniques employed in this thesis. Two of my professors also deserve special thanks: Professor Ravi Ramakrishna, for teaching me algebra and providing guidance throughout my undergraduate career, and Professor Marshall Cohen, who taught me algebraic topology from the simplicial viewpoint. Finally, I would like to thank the Cornell Presidential Research Scholars program, which provided me with funding for the first year of this project.

Chapter 1

Introduction

At the intersection of topology, algebra and combinatorics lies the study of simplicial complexes arising from finite groups. Over the last twenty-five years, there has been a great deal of research in this area, largely stimulated by Quillen's influential paper [18] on the p -group complex. A number of other references in the area are listed in the bibliography [1, 2, 7, 14, 17, 21, 22, 20]. This thesis represents a contribution to the theory, focusing in particular on a new complex (the coset poset) recently introduced by Brown [7].

The coset poset of a group finite G , denoted $\mathcal{C}(G)$, is the partially ordered set containing all cosets of all proper subgroups of G (ordered by inclusion). One forms a simplicial complex on the vertex set $\mathcal{C}(G)$ by declaring each chain

$$x_1H_1 < \cdots < x_nH_n$$

in $\mathcal{C}(G)$ to be a simplex. This complex arose from Brown's analysis of $P(G, s)$, the probabilistic zeta function of a finite group. This function gives the probability that s elements of G , chosen at random, generate G . The theory of Möbius inversion allows one to write $P(G, s)$ in the form

$$P(G, s) = \sum_{H \leq G} \frac{\mu(H, G)}{(G : H)^s},$$

where μ is the Möbius function of the subgroup lattice of G (defined in Section 4.3.2). Written as such, we see that $P(G, s)$ is defined for any complex number $s \in \mathbb{C}$. The starting point of [7] is the observation that $P(G, -1)$ is in fact the Euler characteristic of $\mathcal{C}(G)$.

This result motivates the study of the homotopy type of $\mathcal{C}(G)$. In this thesis, we will mainly be concerned with connectivity. In particular we will investigate the question of simple connectivity, raised in Brown [7, Question 4]. Although we are not able to give a characterization of groups with simply connected coset posets, we do present several results in either direction.

The structure of this thesis is as follows. In Chapter 2, we develop the basic topological theory of simplicial complexes and partially ordered sets. This theory is developed from scratch, assuming only standard topological prerequisites. After describing the geometric realization functors and proving some of their basic properties, we proceed to prove a variety of results on the homotopy type of posets and simplicial complexes. Most of this background material is taken from Björner [3].

In Chapter 3, we introduce the coset poset and prove a variety of new results on its connectivity. Some of these results extend to a more general class of posets we call *atomized posets*. These are essentially posets in which any set of minimal elements “generates” a well-defined element of the poset (or generates the whole poset). In addition to $\mathcal{C}(G)$ this class includes, for example, the poset of proper, non-trivial subgroups of a finite group. In these cases, the notion of generation in the poset coincides with the group-theoretic definition.

The three central methods in Chapter 3 are the introduction of the minimal cover of an atomized poset, the use of Mayer-Vietoris sequences to analyze the homology of $\mathcal{C}(G)$, and the technique (introduced in Brown [7]) of finding a large subposet that can be described as a join. The minimal cover of $\mathcal{C}(G)$ is the simplicial complex with vertex set G and with a simplex for each subset of a (proper) coset of G . Denoting the minimal cover by $\mathcal{M}(G)$, we show that $\mathcal{M}(G) \simeq \mathcal{C}(G)$ (Lemma 3.1.4). For many purposes, $\mathcal{M}(G)$ is easier to analyze, and many of our results are based on the structure of this complex. For example, we prove that any group not generated by k elements has a $(k - 1)$ -connected coset poset (Theorem 3.1.5). If G is a finite 2-generator group this theorem says only that $\mathcal{C}(G)$ is connected, but with additional hypotheses we show that $\mathcal{C}(G)$ is simply connected as well (Theorem 3.1.9).

Using Mayer-Vietoris sequences, we show that certain classes of finite groups have non-simply connected coset posets (Theorems 3.2.1 and 3.2.4). In addition, we give lower bounds on the rank of the first homology of $\mathcal{C}(G)$ for these groups. These results lead to a purely group-theoretic result on the number of split factors in a chief series for certain groups (Corollaries 3.3.3

and 3.3.4).

In the last section of Chapter 3, we consider the coset poset of a group G written as an extension $N \hookrightarrow G \twoheadrightarrow G/N$. As a corollary of Brown's analysis of products [7, Lemma 5], we deduce a simple characterization of those direct products with simply connected coset posets. In addition, we generalize [7, Lemma 5] to the case of a semi-direct product, and note a simple result about general extensions.

In the final chapter, we compute a number of interesting examples. First, we describe a third complex homotopy equivalent to $\mathcal{C}(G)$. This complex immediately gives the homotopy-type of $\mathcal{C}(Q_8)$, where Q_8 denotes the Quaternion group of eight elements. The computation provides a counterexample to the converse of an earlier result. (Note that even for a small group such as Q_8 , the coset poset is too large to admit direct analysis: $\mathcal{C}(Q_8)$ is a two-dimensional complex with eighteen vertices, forty-four edges and twenty-four triangles.)

Next, we turn to finite simple groups. Brown has given a complete description of the homotopy-type of $\mathcal{C}(G)$ for any finite solvable group G , and our results about extensions suggest that the existence of a non-trivial normal subgroup greatly simplifies the coset poset. This motivates the study of simple groups, at the other end of the spectrum. First we calculate the homotopy-type of $\mathcal{C}(A_5)$, giving two simpler proofs of an unpublished result of Shareshian (cited in [7, p. 1009]). The first proof makes use of a theorem from Chapter 3, and the second utilizes 2-transitive actions.

We also consider the coset poset of $PSL_2(\mathbb{F}_7)$. We begin by describing in detail the subgroup lattice of the finite simple groups $PSL_2(\mathbb{F}_p)$, following the analysis given in Burnside [8]. We then show that $\mathcal{C}(PSL_2(\mathbb{F}_7))$ is simply connected and make a few other observations about its homotopy-type.

In the final chapter we suggest several new directions for research and formulate some conjectures on the homotopy type of $\mathcal{C}(G)$.

Chapter 2

Background Material

In this chapter we will introduce the topological definitions and machinery that will be used through this thesis. We will deal mainly with simplicial complexes, and will focus in particular on simplicial complexes arising from finite partially ordered sets (posets).

Throughout this thesis, the word map will mean a morphism in the appropriate category. When the category is not clear from context, we will be more explicit.

2.1 Simplicial Complexes

Definition 2.1.1 *An abstract simplicial complex is a pair $S = (V, \Delta)$ where V is a finite set (called the vertices of S) and Δ is a collection of subsets of V (called the simplices of S) satisfying the following properties:*

1. *If $\sigma_1 \subset \sigma_2$ and $\sigma_2 \in \Delta$, then $\sigma_1 \in \Delta$,*
2. *For all $v \in V$, $\{v\} \in \Delta$.*

In other words, Δ is closed under inclusion and contains all singleton-sets.

For any simplex $\sigma \in \Delta$, let $\dim \sigma = |\sigma| - 1$. If $\dim \sigma = n$ we call σ an n -simplex.

From here on, *simplicial complex* will mean abstract simplicial complex. If $S = (V, \Delta)$ is a simplicial complex, then by condition 2, $V = \bigcup_{\sigma \in \Delta} \sigma$ and

we may suppress V from the notation and denote the complex simply by Δ . When necessary, we will denote the vertex set of Δ by $V(\Delta)$.

If Δ is a simplicial complex and $\Gamma \subset \Delta$ is also a simplicial complex, then we call Γ a *subcomplex* of Δ . For $k = 0, 1, \dots$ we define Δ^k , the k -skeleton of Δ , to be the subcomplex containing all simplices of dimension at most k .

To each simplicial complex we will now associate a topological space, called the *geometric realization* of the complex.

For any finite set X , let $\mathbb{R}^X$ be the $\mathbb{R}$ -vector space with basis X (so $\mathbb{R}^X \cong \mathbb{R}^{|X|}$).

If $t = \sum_{i=1}^n t_i x_i \in \mathbb{R}^X$ ($x_i \in X$), then the *support* of t is the set $\text{supp}(t) = \{x_i : t_i \neq 0\}$. We will sometimes denote the point t simply by $(t_1, \dots, t_n)$.

Definition 2.1.2 *Let Δ be a simplicial complex with vertex set V . Then $|\Delta|$, the geometric realization of Δ , is the subspace of $\mathbb{R}^V$ given by*

$$|\Delta| = \{t = (t_1, \dots, t_n) \in \mathbb{R}^V : t_i \geq 0 \ \forall i, \sum_{i=1}^n t_i = 1 \text{ and } \text{supp}(t) \in \Delta\}.$$

The geometric realization of a non-empty simplex $\sigma \in \Delta$ is the subspace $|\sigma| = \{\sum_{v \in V} t_v v \in |\Delta| : t_i = 0 \text{ if } v \notin \sigma\}$.

The geometric realization of a simplex thus consists of all weighted averages of the vertices of that simplex, and the geometric realization of a complex is the union of the geometric realizations of its simplices. Thus the geometric realizations of 0-, 1-, 2-, and 3-simplicexes are points, lines, triangles and tetrahedrons respectively.

Although the same notation is used for both cardinality and geometric realization, in context it will always be clear what is meant, or else we will be more explicit. Also, we will sometimes refer simply to Δ , and when references are made the topology of Δ this will always mean the topology of $|\Delta|$.

An important class of maps between simplicial complexes are the *simplicial maps*, maps that respect the combinatorial sturcture of the complexes in question.

Definition 2.1.3 *Let Δ and Γ be simplicial complexes. We call a map $f : V(\Delta) \rightarrow V(\Gamma)$ simplicial if for each simplex $\sigma \in \Delta$, $f(\sigma) \in \Gamma$. (Note that f need not be bijective; $\dim f(\sigma) < \dim \sigma$ is allowed.)*

Any simplicial map $f : V(\Delta) \rightarrow V(\Gamma)$ induces a map $|f|$ on geometric realizations, called the geometric realization of f . For $t = \sum_{v \in V(\Delta)} t_v v \in |\Delta|$, we define $|f|(t) = \sum_{v \in V(\Delta)} t_v f(v)$.

A simplicial map $f : V(\Delta) \rightarrow V(\Gamma)$ induces a map $f' : \Delta \rightarrow \Gamma$, and we will usually introduce a simplicial map in the latter form. So a map $f' : \Delta \rightarrow \Gamma$ is simplicial iff it is induced by some simplicial map $f : V(\Delta) \rightarrow V(\Gamma)$.

One important fact about geometric realizations is that the realization of a simplicial map is always continuous. Together with the (easy) fact that $|f \circ g| = |f| \circ |g|$ (i.e. geometric realizations commute with compositions) this will mean that $|\cdot|$ is a *functor* from the category of simplicial complexes and simplicial maps to the category of topological spaces and continuous maps. In order to prove that the geometric realization of a simplicial map is continuous, we will need the following basic result about the topology of a simplicial complex.

Claim 2.1.4 *Let Δ be a simplicial complex. Then a set $U \subset |\Delta|$ is open iff $U \cap |\sigma|$ is open in $|\sigma|$ for each $\sigma \in \Delta$. Also, if X is any topological space, a map $f : |\Delta| \rightarrow X$ is continuous iff its restriction to each simplex $|\sigma|$ is continuous.*

Now we show that the geometric realization of a simplicial map is continuous.

Claim 2.1.5 *Let $f : \Delta \rightarrow \Gamma$ be a simplicial map. Then $|f| : |\Delta| \rightarrow |\Gamma|$ is continuous.*

Proof. By Claim 2.1.4 we simply need to show that the restriction of $|f|$ to a simplex $|\sigma| \subset |\Delta|$ is continuous. Setting $f(\sigma) = \tau$, it suffices to check that the map $\hat{f} : |\sigma| \rightarrow |\tau|$, $\hat{f}(t) = |f|(t)$, is continuous.

Let $V \subset |\tau|$ be open, let $U = \hat{f}^{-1}(V)$ and let $\dim \sigma = n$. For any $x \in U$, we may choose $\epsilon > 0$ s.t. all points of the form $t = f(x) + \sum_{v \in \tau} \epsilon_v v$, with $|\epsilon_v| < \epsilon$ for each v and $t \in \tau$, are in V (this is possible because V is open in τ). Then any point of the form $x' = x + \sum_{u \in \sigma} \frac{\epsilon_u}{n+1} u$, with $|\epsilon_u| < \epsilon$ and with $x' \in |\sigma|$, is in U since $\hat{f}(x') = f(x) + \sum_{u \in \sigma} \frac{\epsilon_u}{n+1} f(u)$, and the coefficient on any particular $v \in \tau$ has absolute value at most $(n+1) \frac{\epsilon}{n+1} = \epsilon$. $\square$

Our next goal will be to give a presentation for the fundamental group of an arbitrary simplicial complex in terms of the combinatorial structure of its 2-skeleton. In order to do this, we will first need to show that for any simplicial complex Δ we have $\pi_1(\Delta) \cong \pi_1(\Delta^2)$. We will in fact prove this latter fact for CW-complexes, and our proof will be a simple application

of the Cellular Approximation Theorem. For standard definitions and facts about CW-complexes, we refer to Hatcher [12].

We denote the n -skeleton of a CW-complex X by X^n , and we call a map $f : X \rightarrow Y$, with X and Y CW-complexes, *cellular* if $f(X^n) \subset Y^n$ for each n .

If Δ is a simplicial complex, then $|\Delta|$ has an obvious CW structure, and the definitions of the skeleta coincide. In addition, the geometric realization of a simplicial map is cellular.

Theorem 2.1.6 (Cellular Approximation Theorem) *Let $f : X \rightarrow Y$ be a map between two CW-complexes. If f is cellular on the n -skeleton of X , ($n = 0, 1, \dots$) then there exists a cellular map $f' : X \rightarrow Y$ s.t. $f \simeq f'$ (rel X^n).*

Corollary 2.1.7 *If Δ is a (connected) simplicial complex, then $\pi_1(\Delta) \cong \pi_1(\Delta^2)$.*

Proof. Fix a basepoint $v \in V(\Delta)$. Consider the map $f : \pi_1(\Delta^2) \rightarrow \pi_1(\Delta)$ induced by the inclusion $\Delta^2 \hookrightarrow \Delta$. We must show that this is a bijection. First, note that each homotopy class of loops has a cellular representative, by Theorem 2.1.6. This immediately means that f is surjective, since any cellular loop is contained in $|\Delta^1| \subset |\Delta^2|$.

Now, in order to prove injectivity we must show that if $\alpha, \beta : I \rightarrow |\Delta^2|$ are two cellular loops based at v , then $\alpha \simeq \beta$ in $|\Delta|$ implies $\alpha \simeq \beta$ in $|\Delta^2|$. But this is immediate from Theorem 2.1.6, since the homotopy in $|\Delta|$ may be pushed down into $|\Delta^2|$ keeping its restrictions α and β fixed. $\square$

Next, we recall that given any presentation $\mathcal{P} = \langle S | R \rangle$ where S is a set of generators and R is a set of relators (words in the symbols of S and their formal inverses) there is a CW-complex $K_{\mathcal{P}}$ whose fundamental group has $\mathcal{P}$ as a presentation. We will describe the construction in the case where S and R are each finite.

This complex is formed by starting with a wedge of circles, one for each element $s \in S$ and then attaching 2-cells for each word $r \in R$. Formally, the 1-skeleton of $K_{\mathcal{P}}$ is the disjoint union of $|S|$ copies of I , (we denote these copies by I_s , $s \in S$) mod the relation $0_s \sim 1_s \sim 0_t \sim 1_t$ for all $s, t \in S$. We let q denote the quotient map.

In order to attach the 2-cells, we must first choose orientations on the 1-cells (the images of the copies of I under q). Given $s \in S$, we choose a

map $f_s : I \rightarrow I_s$, where either $f_s(t) = t$ or $f_s(t) = 1 - t$. Composing f_s with q yields the orientation on the 1-cell $q(I_s)$.

If $r = s_0^{\epsilon_0} \cdots s_n^{\epsilon_n}$ (where $s_i \in S$ and $\epsilon_i = \pm 1$) then the 2-cell corresponding to r is attached via the map $\phi_r : I \rightarrow \bigvee_{s \in S} q(I_s)$, where $\phi_r|_{[\frac{i}{n+1}, \frac{i+1}{n+1}]}$ is just a linear shift of the map f_l . (To be more specific, say $0 \leq a < b \leq 1$ and let $L_{[a,b]} : [a,b] \rightarrow I$ be the homeomorphism given by $L_{[a,b]}(x) = \frac{x-a}{b-a}$. Then

$$\phi_r|_{[\frac{i}{n+1}, \frac{i+1}{n+1}]} = f_l \circ L_{[\frac{i}{n+1}, \frac{i+1}{n+1}]}.)$$

Note that due to the choice of orientations, there is some non-uniqueness in the definition of $K_{\mathcal{P}}$. We will ignore this and for any choice of orientations we denote the complex obtained by $K_{\mathcal{P}}$.

From this definition, Van Kampen's Theorem can be used to show that $\mathcal{P}$ is indeed a presentation for $\pi_1(K_{\mathcal{P}})$.

Before we describe the presentation for π_1 of an arbitrary simplicial complex, we need the following lemma.

Lemma 2.1.8 *Let Δ be a simplicial complex, and let $K \subset \Delta$ be a contractible subcomplex. Then $\Delta \simeq \Delta/K$.*

Proof. Since K is contractible, there is a map $C : K \times I \rightarrow K$ satisfying $C_0 = id_K$ and $C_1(K) = k_0$ for some $k_0 \in K$. By the Homotopy Extension Property, this map extends to a map $F : \Delta \times I \rightarrow \Delta$, with $F_0 = id_{\Delta}$. The map F_1 factors through Δ/K , giving a map $\phi : \Delta/K \rightarrow \Delta$.

We claim that ϕ is a homotopy inverse to the quotient map $\pi : \Delta \rightarrow \Delta/K$. First, $\phi\pi = F_1 \simeq F_0 = id_{\Delta}$. To show that $\pi\phi \simeq id_{\Delta/K}$, consider the map $\pi F : \Delta \times I \rightarrow \Delta/K$. This map factors through $\Delta/K \times I$, yielding a map $\Phi : \Delta/K \times I \rightarrow \Delta/K$. It is now easy to check that $\Phi_0 = id_{\Delta/K}$ and $\Phi_1 = \pi\phi$. $\square$

The map F in the above proof may be described explicitly; see [4].

Theorem 2.1.9 *Let Δ be a simplicial complex, and let $T \subset \Delta^1$ be a maximal tree (i.e. a spanning tree).*

Then $\pi_1(\Delta)$ has a presentation with a generator for each (ordered) edge (u, v) with $\{u, v\} \in \Delta^1$, and with the following relations:

1. $(u, v) = 1$ if $\{u, v\} \in T$,
2. $(u, v)(v, u) = 1$ if $\{u, v\} \in \Delta^1$,

3. $(u, v)(v, w)(w, u) = 1$ if $\{u, v, w\} \in \Delta^2$.

Proof. We begin by simplifying the presentation given above.

For each 1-simplex $\{u, v\} \in \Delta^1 - T$, choose a (combinatorial) orientation (either (u, v) or (v, u)) and similarly for each 2-simplex. (These orientations are just orderings of the vertices and we do not require them to agree with one another in any way.) Let $\mathcal{O}_1$ be the set of chosen oriented edges and let $\mathcal{O}_2$ be the set of chosen oriented 2-simplices.

Let F denote the free group generated by $\mathcal{O}_1$. First, if (u, v) is the chosen orientation for an edge $\{u, v\} \in \Delta$, then we let (v, u) denote $(u, v)^{-1}$ (the inverse of (u, v) in F). Also, for $\{u, v\} \in T$, we let $(u, v) \equiv (v, u) \equiv 1$ (i.e. (u, v) and (v, u) are formal symbols representing the identity in F).

Then the following presentation $\mathcal{P}$ clearly realizes the same group as the presentation given in the theorem:

$$\mathcal{P} = \langle \mathcal{O}_1 | (u, v)(v, w)(w, u) = 1 \text{ if } (u, v, w) \in \mathcal{O}_2 \rangle.$$

By Lemmas 2.1.7 and 2.1.8 we have

$$\pi_1(\Delta) \cong \pi_1(\Delta^2) \cong \pi_1(\Delta^2/T).$$

Now, Δ^2/T is a CW-complex whose cells are the images of the simplices in Δ^2 . We claim that in fact, Δ^2/T is (homeomorphic to) the standard complex for the presentation $\mathcal{P}$.

Let $q : \Delta^2 \rightarrow \Delta^2/T$ be the quotient map. Then Δ^2/T has a unique vertex $v_0 = q(V(\Delta))$ and a 1-cell for each generator in $\mathcal{O}_1$. In defining the standard complex for $\mathcal{P}$, it is necessary to choose an orientation on the 1-cells. We will choose to orient the 1-cells of Δ^2/T in accordance with the chosen orientations on the edges of Δ^2 , as follows. Say $(u, v) \in \mathcal{O}_1$. Then let $f_{(u, v)} : I \rightarrow \Delta$ be the map $f_{(u, v)}(t) = (1 - t)u + tv$. It is clear that $f_{(u, v)}(I) = |\{u, v\}|$. Now the orientation on the 1-cell corresponding to $\{u, v\}$ is simply $q \circ f_{(u, v)}$, and this agrees with the definition of $K_{\mathcal{P}}$.

Now consider a 2-cell $\tau = q(|\{u, v, w\}|)$, where $\{u, v, w\} \in \Delta$. Assume that (u, v, w) is the chosen orientation on this simplex. Then the attaching map for c is just (linear shifts of) the orienting map $f_{(u, v)}$ followed by $f_{(v, w)}$ followed by $f_{(w, u)}$, just as in the standard complex $K_{\mathcal{P}}$. $\square$

2.2 Homotopy Theory for Posets

In this section, we will describe a functor from the category of posets (partially ordered sets) to the category of simplicial complexes. This will allow us to apply topological concepts to posets, and in particular we will be able to talk about the homotopy type of a poset. After introducing this functor, we will prove several results involving homotopy equivalence and homotopy type of posets.

Definition 2.2.1 *A poset is a set P together with a subset $R \subset P \times P$ (called a partial order on P) satisfying the following properties:*

1. $(x, x) \in R$ for all $x \in P$ (reflexive).
2. If $(x, y) \in R$ and $(y, z) \in R$ then $(x, z) \in R$ (transitive).
3. If $(x, y) \in R$ and $(y, x) \in R$, then $x = y$ (anti-symmetric).

When $(x, y) \in R$, we write $x \leq y$. If in addition $x \neq y$, we write $x < y$. We call such elements x and y comparable.

If $S \subset P$ has a least upper bound $j \in P$, we call j the join of S and write $m = \vee S$. Similarly, if S has a greatest lower bound $m \in P$, we call m the meet of P and write $m = \wedge S$.

From now on, we assume that all posets are finite. Given a poset P , the *order complex* of P (denoted $\Delta(P)$) is the simplicial complex whose vertices are the elements of P and whose simplices are the *chains* of P , i.e. the sequences $x_1 < x_2 < \cdots < x_n$. It is clear that $\Delta(P)$ is a simplicial complex.

If P and Q are posets, a map $f : P \rightarrow Q$ is called *order-preserving* if $x \leq y \implies f(x) \leq f(y)$. Note that the composition of two order-preserving maps is order-preserving. An order-preserving map of posets induces a simplicial map between order complexes in the obvious manner. For simplicity, we use the same symbol to denote the map $f : P \rightarrow Q$ and the induced map $\Delta(P) \rightarrow \Delta(Q)$, and we let $|P|$ denote $|\Delta(P)|$ (this will be called the geometric realization of P). Note that order-reversing maps also induce simplicial maps on order complexes. [To keep the relationship between P and $\Delta(P)$ functorial, one may think of an order reversing map $f : P \rightarrow Q$ as an order-preserving map to the dual poset Q^* , in which $x \leq_{Q^*} y \iff y \leq_Q x$. Then simply note that $\Delta(Q) = \Delta(Q^*)$.]

Definition 2.2.2 *Let P and Q be posets. We say P and Q are homotopy equivalent ($P \simeq Q$) if the geometric realizations of their order complexes are homotopy equivalent.*

We can now state an important theorem of Quillen [18, Proposition 1.6] giving conditions under which an order-preserving map of posets induces a homotopy equivalence of order complexes.

Theorem 2.2.3 (Quillen, 1978) *Let P and Q be posets and let $f : P \rightarrow Q$ be an order-preserving map. For $x \in Q$, let $Q_{\geq x}$ be the set of all elements greater than or equal to x . If $|f^{-1}(Q_{\geq x})|$ is contractible for each $x \in Q$, then $|f|$ is a homotopy equivalence from $|P| \rightarrow |Q|$.*

Note that any subset of a poset is again a poset, and the order complex of a subposet is a full subcomplex of the order complex.

The proof of Quillen's Theorem presented here comes from [3] and is an application of the Carrier Lemma. In order to prove this lemma, we will first need a basic fact about maps into contractible spaces.

Given a topological space X , the *cone* on X is the space

$$C(X) = (X \times I) / X \times \{1\}.$$

Lemma 2.2.4 *Let X and Y be topological spaces, and assume that Y is contractible. Then any map $f : X \rightarrow Y$ can be extended to a map $\hat{f} : C(X) \rightarrow Y$ satisfying $\hat{f} \circ q(x, 0) = f(x)$ for each $x \in X$ (here q denotes the quotient map $X \times I \rightarrow C(X)$).*

Proof. Since Y is contractible, there is a map $H : Y \times I \rightarrow Y$ and a point $y_0 \in Y$ s.t. $H(y, 0) = y$ and $H(y, 1) = y_0$ for each $y \in Y$. Letting $f \times id_I : X \times I \rightarrow Y \times I$ denote the map $(x, t) \mapsto (f(x), t)$, the composition $H \circ (f \times id_I) : X \times I \rightarrow Y$ factors through the quotient map q and gives the desired extension of f . $\square$

Let Δ be a simplicial complex and let X be any topological space. We call a function $K : \Delta \rightarrow 2^X$ (the power set of X) a *contractible system* if

1. $\sigma \subset \tau \implies K(\sigma) \subset K(\tau)$
2. $K(\sigma)$ is contractible for each simplex σ .

Also, if $f : |\Delta| \rightarrow X$ is a continuous map, we say f is *carried* by K if $f(|\sigma|) \subset K(\sigma)$ for each $\sigma \in \Delta$. The following lemma shows that maps carried by a particular system K exist and are unique up to homotopy.

Lemma 2.2.5 (Carrier Lemma) *Let Δ be a simplicial complex, X a topological space and K a contractible system. Then there exists a continuous map $f : |\Delta| \rightarrow X$ carried by K , and if g is also carried by K then $f \simeq g$.*

Proof. First we construct a map f carried by K . We will construct f inductively, beginning with the vertices of Δ . If $v \in V(\Delta)$, let $f(v)$ be any point in $K(v)$. Now assume that we have a map $f : \Delta^k \rightarrow X$ carried by K , and let σ be a $k+1$ -simplex. Clearly, $|\sigma| \cong C(\sigma^k)$ and $f(\sigma^k) \subset K(\sigma)$, which is contractible. So Lemma 2.2.4 applies, and $f|_{\sigma^k}$ extends to a map $|\sigma| \rightarrow K(\sigma)$. Since σ is finite, we may extend the map f to the entire $k+1$ -skeleton of Δ , and repeating this process will eventually give a map from Δ to X carried by K .

Next, say $g : |\Delta| \rightarrow X$ is also carried by K . We need to construct a homotopy $F : |\Delta| \times I \rightarrow X$ with $F_0 = f$ and $F_1 = g$. This can be done in a manner similar to the construction of f above. Again we define F inductively on successive skeleta of Δ . Note that it suffices to check the continuity of F on the set of $|\sigma| \times I$, $\sigma \in \Delta$ (these sets are closed in the product topology, and cover $\Delta \times I$).

If v is a vertex of Δ , then $f(v), g(v) \in C(v)$ and since $K(v)$ is contractible, it is path-connected, i.e. there is a map $H : I \rightarrow K(v)$ s.t. $H(0) = f(v)$ and $H(1) = g(v)$. We define $F|_{\{v\} \times I} = H \circ p$, where p is the projection $\{v\} \times I \rightarrow I$. Now, if F has been appropriately defined on $\Delta^k \times I$ and σ is a $k+1$ -simplex of Δ , then $F|_{|\sigma^k| \times I}$ extends as before to a map of $C(|\sigma^k| \times I) \cong |\sigma| \times I$. $\square$

Proof of Theorem 2.2.3. Let P and Q be posets, and let $f : P \rightarrow Q$ be an order-preserving map s.t. for each $x \in Q$, $f^{-1}(Q_{\geq x})$ is contractible. For any simplex $\sigma \in \Delta(Q)$, define $K(\sigma) = |f^{-1}(Q_{\geq \min(\sigma)})| \subset |P|$ (note that σ is a chain in Q and thus has a minimum element). K is a contractible system by hypothesis, and Lemma 2.2.5 gives us a map $g : |Q| \rightarrow |P|$ carried by K . We claim that g is a homotopy inverse to $|f|$ (which from here on we denote simply by f).

First we show that $g \circ f \simeq id_P$. Consider the contractible system $K' : P \rightarrow 2^{|P|}$ given by $K'(\tau) = |f^{-1}(Q_{\geq \min f(\tau)})|$. We claim that K' carries both $g \circ f$ and id_P . Clearly K' carries id_P , since if σ is a chain in P , then $f(\sigma)$ lies above $\min f(\sigma)$ and hence $\sigma \subset K(\sigma)$. Also, if $\tau \in \Delta(P)$ then

$$g \circ f(\tau) \subset K(f(\tau)) = |f^{-1}(Q_{\geq \min(f(\tau))})| = K'(\tau).$$

Lemma 2.2.5 now shows that $g \circ f \simeq id_P$.

The proof that $f \circ g \simeq id_Q$ is similar. We define the contractible system $K'' : Q \rightarrow 2^{|Q|}$ by $K''(\sigma) = |Q_{\geq \min(\sigma)}|$ and as before, it is clear that K'' carries id_Q . K'' also carries $f \circ g$, which can be checked as follows:

$$f \circ g(\sigma) \subset f(K(\sigma)) = f(|f^{-1}(Q_{\geq \min(\sigma)})|) \subset |Q_{\geq \min(\sigma)}| = K''(\sigma).$$

□

Our next goal is to prove the Nerve Theorem, which will be useful in our study of the coset poset. Before proving this theorem, need a technical result about simplicial complexes.

If Δ is a simplicial complex, the *face poset* of Δ is the poset $P(\Delta)$ whose elements are the simplices of Δ , ordered by inclusion. The simplicial complex $sd(\Delta) = \Delta(P(\Delta))$ is the *first barycentric subdivision* of Δ and we have:

Proposition 2.2.6 *If Δ is a simplicial complex, then $|\Delta|$ is homeomorphic to $|sd(\Delta)|$.*

Proof. We will define an explicit homeomorphism $h : |sd(\Delta)| \rightarrow |\Delta|$. First we define h on the vertices of $|sd(\Delta)|$, again using the convention that the standard basis vectors of the vector spaces containing $|\Delta|$ and $|sd(\Delta)|$ are identified with the vertices of these complexes.

For any $\sigma = \{v_1, \dots, v_n\} \in V(sd(\Delta)) = \Delta$, we define

$$h(\sigma) := \frac{1}{\dim \sigma + 1} \sum_{i=1}^n v_i.$$

(The point $h(\sigma)$ is called the *barycenter* of σ .) Now we extend h by linearity (so we may view h as a linear transformation of vector spaces), so that for an arbitrary point $t = \sum_{i=1}^k t_i \sigma_i \in \mathbb{R}^\Delta$,

$$h(t) = \sum_{i=1}^k t_i h(\sigma_i).$$

Next we show that $h(|sd(\Delta)|) \subset |\Delta|$. Consider an arbitrary point $t \in |sd(\Delta)|$. We may write $t = \sum_{i=1}^k t_i \sigma_i$, and the simplices σ_i must form a simplex in $sd(\Delta)$, i.e. they must form a chain in the face poset $P(\Delta)$. Without loss of generality we may assume that $\sigma_1 \subset \sigma_2 \subset \dots \subset \sigma_k$. Let

$\sigma_k = \{v_1, \dots, v_n\}$, and let $\sigma_i = \{v_1, \dots, v_{n_i}\}$ (so the n_i 's form an increasing sequence).

Then we have

$$h(t) = \sum_{i=1}^k t_i h(\sigma_i) = \sum_{i=1}^k \frac{t_i}{n_i} \sum_{j=1}^{n_i} v_j,$$

and thus the total sum of the coefficients of $h(t)$ is

$$\sum_{i=1}^k \frac{t_i}{n_i} n_i = \sum_{i=1}^k t_i = 1.$$

Also, all the terms above are positive, so the coefficient of any particular v_i is positive. Finally, the support of $h(t)$ is contained in σ_k and is thus a simplex of Δ . So $h(t) \in |\Delta|$ as desired.

Since h is continuous on $\mathbb{R}^\Delta$, its restriction to $|sd(\Delta)|$ is a continuous map from $|sd(\Delta)|$ to $h(|sd(\Delta)|)$. Thus it remains to show that h is in fact a bijection between $|sd(\Delta)|$ and $|\Delta|$. This will complete the proof, because a continuous bijection of compact hausdorff spaces is a homeomorphism.

First we show that h is injective. We will actually prove a bit more than this.

For any poset P , let

$$|\widehat{P}| = \left\{ \sum_{i=1}^n \lambda_i p_i \in \mathbb{R}^P : \lambda_i > 0, \ 0 < \sum_{i=1}^n \lambda_i \leq 1 \text{ and } p_1 < \dots < p_n \right\}.$$

If $x \in |\widehat{P}|$, we may write x uniquely as $\sum_{i=1}^n \lambda_i p_i$ (with $p_1 < \dots < p_n$), and we define $\nu(x) = n$ and $\lambda(x) = \sum_{i=1}^n \lambda_i$.

Claim: The restriction of h to $|\widehat{sd(\Delta)}|$ is injective.

Proof of Claim. Say $s, t \in |\widehat{sd(\Delta)}|$ and $h(s) = h(t)$. We will show, by induction on $m(s, t) := \min(\nu(s), \nu(t))$, that $s = t$.

First, say $m(s, t) = 1$. Assume that $\nu(s) = 1$, so that $s = \lambda(s)\sigma$ for some $\sigma \in \Delta$. Letting $\sigma = \{v_1, \dots, v_p\}$, we have

$$h(s) = \frac{\lambda(s)}{p} \sum_{i=1}^p v_p.$$

Next, let $t = \sum_{i=1}^l t_i \tau_i$, with $\tau_1 < \dots < \tau_l$ (so $l = \nu(\tau)$). Also, let $\tau_i = \{w_1, \dots, w_{q_i}\}$. Then we have

$$h(t) = \sum_{i=1}^l \frac{t_i}{q_i} \sum_{j=1}^{q_i} w_j,$$

and we see that the support of $h(t)$ is exactly $\{w_1, \dots, w_{q_l}\}$ (since $t_i > 0$ for $i = 1, \dots, l$).

Thus, since $h(s) = h(t)$, $\{w_1, \dots, w_{q_l}\} = \sigma$, and (by renumbering the v_i) we may assume $w_i = v_i$. Equating the coefficients of $v_p = w_{q_l}$ in the expressions for $h(s)$ and $h(t)$ gives

$$\frac{\lambda(s)}{p} = \frac{t_l}{q_l} = \frac{t_l}{p}$$

and thus $t_l = \lambda(s)$. So the l -th term of the expression for $h(t)$ is in fact the entire expression (because $t_l = \lambda(s) = \lambda(h(s)) = \lambda(h(t))$), so $l = 1$ and $t = \lambda(s)\sigma = s$.

Now assume the Claim for any pair of points s', t' (satisfying the hypotheses) with $m(s, t) = n$ ($n \geq 1$), and consider points s, t with $m(s, t) = n + 1$.

By the same reasoning as above, if $s = \sum_{i=1}^k s_i \sigma_i$ and $t = \sum_{i=1}^l t_i \tau_i$ are in $\widehat{sd(\Delta)}$ (where $\sigma_1 \subset \dots \subset \sigma_k$ and $\tau_1 \subset \dots \subset \tau_k$) then we have $\sigma_k = \tau_l$. Let this simplex be $\{v_1, \dots, v_p\}$, and let $\sigma_i = \{v_1, \dots, v_{p_i}\}$, $\tau_i = \{v_1, \dots, v_{q_i}\}$. Then, since $h(s) = h(t)$, we have

$$\sum_{i=1}^k \frac{s_i}{p_i} \sum_{j=1}^{p_i} v_j = \sum_{i=1}^l \frac{t_i}{q_i} \sum_{j=1}^{q_i} v_j,$$

and by equating the coefficients of $v_p = v_{p_k} = v_{q_l}$ we see that $\frac{s_k}{p_k} = \frac{t_l}{q_k}$, so $s_k = t_l$ (since $p_k = q_k = p$) and thus

$$h\left(\sum_{i=1}^{k-1} s_i \sigma_i\right) = h\left(\sum_{i=1}^{l-1} t_i \tau_i\right).$$

The induction hypothesis applies and gives us that $\sum_{i=1}^{k-1} s_i \sigma_i = \sum_{i=1}^{l-1} t_i \tau_i$. Adding back in the last terms gives $s = t$, completing the proof of the claim. $\square$

Proof of Proposition 2.2.6 (Continued) To complete the proof, we must show that h is surjective.

Consider an arbitrary point $\sum_{i=1}^n a_i v_i \in |\Delta|$, and assume that $a_1 \geq a_2 \geq \dots \geq a_n$.

Let $t = \sum_{i=1}^n t_i \sigma_i$, where $\sigma_i = \{v_1, \dots, v_i\}$. Then

$$h(t) = \sum_{i=1}^n t_i h(\sigma_i) = \sum_{i=1}^n \frac{t_i}{i} \sum_{j=1}^i v_j = \sum_{i=1}^n v_i \sum_{j=i}^n \frac{t_j}{j},$$

and we want to choose $t_1, \dots, t_n$ s.t.

$$\sum_{j=i}^n \frac{t_j}{j} = a_i \quad (2.1)$$

(and s.t. $t \in |sd(\Delta)|$). Note that we may choose $t_i = 0$ for some i .

Equation 2.1 gives us $\frac{t_n}{n} = a_n$, i.e. $t_n = na_n$.

Next, we have $t_{n-1} = (n-1)(a_{n-1} - \frac{t_n}{n}) = (n-1)(a_{n-1} - a_n)$, and assuming inductively that $t_j = j(a_j - a_{j+1})$ for $j > i$ (and with the convention that $a_{n+1} = 0$) we see that

$$\begin{aligned} t_i &= i \left(a_i - \frac{t_n}{n} - \frac{t_{n-1}}{n-1} - \dots - \frac{t_{i+1}}{i+1} \right) \\ &= i(a_i - a_n - (a_{n-1} - a_n) - (a_{n-2} - a_{n-1}) - \dots \\ &\quad - (a_{i+2} - a_{i+3}) - (a_{i+1} - a_{i+2})) \\ &= i(a_i - a_{i+1}). \end{aligned}$$

Since we have assumed that $a_1 \geq \dots \geq a_n$, all of the coefficients determined above are non-negative. Thus to show that $t \in |sd(\Delta)|$, it remains only to check that their sum is exactly 1. We have:

$$\begin{aligned} \sum_{i=1}^n t_i &= \sum_{i=1}^n i(a_i - a_{i+1}) \\ &= (a_1 - a_2) + 2(a_2 - a_3) + \dots \\ &\quad + (n-1)(a_{n-1} - a_n) + na_n \\ &= \sum_{i=1}^n a_i = 1, \end{aligned}$$

the last step following from the fact that $\sum_{i=1}^n a_i v_i \in |\Delta|$. $\square$

Given a finite collection $\{V_i\}_{i \in I}$ of subsets of a set S , we define the *nerve* of V_i to be the simplicial complex $\mathcal{N}(\{V_i\})$ with vertices I and with a simplex for each set $J \subset I$ s.t. $\bigcap_{j \in J} V_j \neq \emptyset$.

Theorem 2.2.7 (Nerve Theorem) *Let Δ be a simplicial complex and let $\{\Gamma_i\}_{i \in I}$ be a collection of subcomplexes. If $\{\Gamma_i\}$ satisfies the following two properties, then $\Delta \simeq \mathcal{N}(\{\Gamma_i\})$:*

1. $\bigcup_{i \in I} \Gamma_i = \Delta$.
2. For any $J \subset I$, the intersection $\bigcap_{j \in J} \Gamma_j$ is contractible.

Proof. By Proposition 2.2.6, it suffices to show that the subdivisions (or face posets) of these complexes are homotopy equivalent. Letting $P(\Delta)^*$ denote the dual poset to $P(\Delta)$ (see the remark after Definition 2.2.1), we have $\Delta(P(\Delta)^*) = sd(P(\Delta))$. We will apply Quillen's Theorem (Theorem 2.2.3) to show that $P(\Delta)^* \simeq P(\mathcal{N})$, where $\mathcal{N} = \mathcal{N}(\{\Gamma_i\}_{i \in I})$.

Consider the map $f : P(\Delta)^* \rightarrow P(\mathcal{N})$ induced by

$$f(\sigma) = \{i \in I : \sigma \in \Gamma_i\},$$

where $\sigma \in V(P(\Delta)) = \Delta$. [Note that $f(\sigma)$ is a simplex of $\mathcal{N}$ (and hence an element of $P(\mathcal{N})$) because $\sigma \in \bigcap_{i \in f(\sigma)} \Gamma_i$.] Now, f is clearly order preserving (since we are using the dual poset $P(\Delta)^*$) and for any $J \subset I$ we have

$$\begin{aligned} f^{-1}(P(\mathcal{N})_{\geq J}) &= \{\sigma \in \Delta : J \subset f(\sigma)\} \\ &= \{\sigma \in \Delta : \sigma \in \bigcap_{j \in J} \Gamma_j\} \\ &= \bigcap_{j \in J} \Gamma_j, \end{aligned}$$

which is contractible by assumption. Quillen's Theorem shows that $|f|$ is a homotopy equivalence, completing the proof. $\square$

In light of Quillen's Theorem and the Nerve Theorem, contractible posets and complexes are of great importance. Our next goal is to give a simple but useful condition under which a poset is contractible. Although the result

could be proven by describing an explicit map, the following approach is more combinatorial in nature and will lead us through a number of other interesting results. In particular, Corollary 2.2.13 will be of use to us in Section 3.4.

Definition 2.2.8 *Let P and Q be posets. We define the product of P and Q to be the set $P \times Q$ together with the ordering $(x, y) \leq (v, w) \iff x \leq v$ and $y \leq w$.*

It is easily checked that this is a partial order on $P \times Q$.

Lemma 2.2.9 *For any finite posets P and Q , we have $|P \times Q| \cong |P| \times |Q|$.*

Proof. The projections $P \times Q \rightarrow P$ and $P \times Q \rightarrow Q$ induce a map $f : |P \times Q| \rightarrow |P| \times |Q|$, with

$$f \left(\sum_{i=1}^n \lambda_i(p_i, q_i) \right) = \left(\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i \right).$$

Note that we may also consider f to be the restriction to $|P \times Q|$ of the map $\mathbb{R}^{P \times Q} \rightarrow \mathbb{R}^P \oplus \mathbb{R}^Q$ sending $(p, q) \in \mathbb{R}^{P \times Q}$ to $(p, q) \in \mathbb{R}^P \oplus \mathbb{R}^Q$.

Since the spaces in question are compact Hausdorff, it suffices to prove that the map f is a bijection. First we prove that f is surjective.

Claim: For any point $(x, y) \in |P| \times |Q|$, $\exists \lambda_1, \dots, \lambda_n \in (0, 1]$, $p_1 \leq \dots \leq p_n \in P$ and $q_1 \leq \dots \leq q_n \in Q$ s.t.

$$(x, y) = \left(\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i \right).$$

Assuming the Claim, we see that f is surjective because if $(x, y) \in |P| \times |Q|$ and $(x, y) = (\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i)$ (where this expression satisfies the conditions in the Claim) then we have $\sum_{i=1}^n \lambda_i(p_i, q_i) \in |P \times Q|$ and

$$f \left(\sum_{i=1}^n \lambda_i(p_i, q_i) \right) = \left(\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i \right) = (x, y).$$

To prove the Claim, we will actually verify a stronger statement.

Claim 2: For any point $(x, y) \in \widehat{|P|} \times \widehat{|Q|}$ with $\lambda(x) = \lambda(y)$ (see the proof of Proposition 2.2.6 for an explanation of this notation), $\exists \lambda_1, \dots, \lambda_n \in (0, 1]$, $p_1 \leq \dots \leq p_n \in P$ and $q_1 \leq \dots \leq q_n \in Q$ s.t.

$$(x, y) = \left(\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i \right).$$

Proof. We proceed by induction on $\nu(x) + \nu(y) =: \nu(x, y)$. If $\nu = 2$ there is nothing to prove, so we may assume the claim for all points (x, y) with $\nu(x, y) \leq r$ (and $\lambda(x) = \lambda(y)$) and consider a point (x, y) with $\nu(x, y) = r+1$. Writing

$$(x, y) = \left(\sum_{i=1}^k t_i p_i, \sum_{i=1}^m s_i q_i \right)$$

with $p_1 < \dots < p_n$ and $q_1 < \dots < q_n$, we may assume WLOG that $t_1 \leq s_1$. We now have

$$(x, y) - (t_1 p_1, t_1 q_1) = \left(\sum_{i=2}^k t_i p_i, (s_1 - t_1) q_1 + \sum_{i=2}^m s_i q_i \right)$$

and it is easy to check that the right-hand side satisfies the induction hypothesis. The claim now follows easily. $\square$

We must now prove that f is injective. Say $x, y \in \widehat{|P \times Q|}$. Then we may write x and y uniquely in the form

$$x = \sum_{i=1}^n \lambda_i (p_i, q_i), \quad y = \sum_{i=1}^m \gamma_i (p'_i, q'_i),$$

with $\lambda_i, \gamma_i > 0$, $(p_1, q_1) < \dots < (p_n, q_n)$ and $(p'_1, q'_1) < \dots < (p'_m, q'_m)$.

Assume that $f(x) = f(y)$, i.e. that

$$\left(\sum_{i=1}^n \lambda_i p_i, \sum_{i=1}^n \lambda_i q_i \right) = \left(\sum_{i=1}^m \gamma_i p'_i, \sum_{i=1}^m \gamma_i q'_i \right).$$

Note that $\sum_{i=1}^n \lambda_i p_i = \sum_{i=1}^m \gamma_i p'_i \implies \{p_1 \dots p_n\} = \{p'_1, \dots, p'_m\} \implies p_1 = p'_1$, and similarly $q_1 = q'_1$.

We will show that $\lambda_1 = \gamma_1$. Since $(p_1, q_1) < (p_2, q_2)$, we may assume WLOG that $p_1 < p_2$. Writing $f(x) = f(y)$ in terms of the standard basis $\{(p, 0)\}_{p \in P} \cup \{(0, q)\}_{q \in Q}$ for $\mathbb{R}^P \oplus \mathbb{R}^Q$, we see that

$$\lambda_1 = \sum_{i: p'_i = p'_1} \gamma_i \geq \gamma_1.$$

If $p'_1 \neq p'_2$, then $\lambda_1 = \sum_{i: p'_i = p'_1} \gamma_i = \gamma_1$ (since $p'_1 < p'_2 \leq \dots \leq p'_m$), and we are done. So assume $p'_1 = p'_2$. Then $q'_1 < q'_2$, and the coefficient on $(0, q'_1) = (0, q_1)$ in $f(x) = f(y)$ is

$$\gamma_1 = \sum_{i: q_i = q_1} \lambda_i \geq \lambda_1.$$

But earlier we saw that $\gamma_1 \leq \lambda_1$, so we must have $\lambda_1 = \gamma_1$. Repeating this process will eventually show that $x = y$. $\square$

For an alternate proof of the above result, see Eilenberg and Steenrod [10].

Lemma 2.2.10 *Let P and Q be posets, and let $f, g : P \rightarrow Q$ be order-preserving maps. If $f(x) \leq g(x)$ for each $x \in P$, then $|f| \simeq |g|$. (We write $f \leq g$ when the above conditions hold.)*

Proof. Let $\mathcal{I} = \{0, 1\}$, with $0 < 1$ (so $|\mathcal{I}|$ is homeomorphic to the unit interval I). Then since $f \leq g$, these maps induce an order-preserving map h on $P \times \mathcal{I}$, with $h(x, 0) = f(x)$, $h(x, 1) = g(x)$. Now $|h|$ is a map from $|P \times \mathcal{I}| \cong |P| \times I \rightarrow |Q|$, and is thus a homotopy from f to g . $\square$

Claim 2.2.11 *If a poset P has a minimum element, then P is contractible.*

Proof. Let p_0 be the minimum element of P . If $f_{p_0} : P \rightarrow P$ is the constant map to p_0 , we have $f_{p_0} \leq id_P$ and the lemma shows that P is contractible. $\square$

Claim 2.2.11 (as well as the generalization below) will be used implicitly throughout this thesis. Using this result, we can extend Lemma 2.2.10. Our proof is taken from Björner [4].

Definition 2.2.12 *Let f and g be order preserving maps from a poset P to a poset Q . If $f(x)$ and $g(x)$ are comparable for all $x \in P$, then we call f and g order-homotopic.*

Corollary 2.2.13 *If f and g are order-homotopic maps from a poset P to a poset Q , then $|f| \simeq |g|$.*

Proof. We will apply the Carrier Lemma. Let $C : \Delta(P) \rightarrow 2^{|Q|}$ be the map $C(\sigma) = f(\sigma) \cup g(\sigma)$, $\sigma \in \Delta(P)$. Note that $C(\sigma)$ always has a minimum element, since $\min f(\sigma) = f(\min \sigma)$ and $\min g(\sigma) = g(\min \sigma)$ are comparable. So C is a contractible system, and since it carries both $|f|$ and $|g|$ we have $|f| \simeq |g|$. $\square$

Using Corollary 2.2.13 we can generalize Lemma 2.2.11.

Definition 2.2.14 *Let P be a poset with an element p_0 which is comparable to each $x \in P$. Then we call P a cone on p_0 , or simply a cone.*

Corollary 2.2.15 *Any cone P is contractible.*

Proof. Let P be a cone on some $p_0 \in P$. Let $f_{p_0} : P \rightarrow P$ be the constant map to p_0 . Then for any $x \in P$, $f_{p_0}(x) = p_0$ is comparable to $\text{id}_P(x) = x$. So f_{p_0} and id are order-homotopic and hence P is contractible. $\square$

Remark 2.2.16 *Cones in the above sense are easily seen to be topological cones at the point p_0 (in the sense of Lemma 2.2.4). This gives another proof of contractibility.*

Chapter 3

Connectivity of the Coset Poset

In this chapter we introduce the coset poset of a finite group and study its homotopy type, using the methods of Chapter 2. We will focus mainly on the connectivity of the coset poset, and in particular on its fundamental group.

Let G be a finite group. The coset poset of G , denoted $\mathcal{C}(G)$, is the poset consisting of all left cosets of all proper subgroups of G , ordered by inclusion (so we allow cosets of the trivial subgroup, but do not allow G itself). Note that the choice of left cosets rather than right cosets is irrelevant, since every left coset is a right coset, and vice versa: $xH = (xHx^{-1})x$. The coset poset was introduced in Brown [7]. This thesis is the result of an attempt to answer a question from [7], asking, “For what groups G is $\mathcal{C}(G)$ simply connected?”

In this chapter, we will present several results describing conditions under which the coset poset is or is not simply connected.

We will also need to consider the subgroup poset of a finite group G . This poset consists of all proper, non-trivial subgroups of G , ordered by inclusion, and is denoted by $S(G)$. We will often use $\mathcal{C}(G)$ and $S(G)$ to denote not only the posets but also the corresponding simplicial complexes and topological spaces.

3.1 Conditions for Simple Connectivity

In this section we will describe two results giving conditions under which the coset poset of a finite group G is simply connected (or k -connected for

higher k). Our results actually extend to a wider class of posets, which we will introduce after discussing some topological preliminaries.

Definition 3.1.1 *Let X be a topological space. We say X is n -connected, $n = 0, 1, 2, \dots$, if for $0 \leq k \leq n$ each map $f : S^k \rightarrow X$ extends to a map $\hat{f} : B^{k+1} \rightarrow X$. We say that X is simply connected if X is 1-connected.*

Note that if X is path-connected, then X is k -connected iff the homotopy group $\pi_k(X)$ is trivial (see Hatcher [12]). Also, 0-connectivity is the same as path-connectivity.

Lemma 3.1.2 *Let Δ be a simplicial complex and say there exists an n -connected simplicial complex Γ with $\Delta^{n+1} = \Gamma^{n+1}$. Then Δ is n -connected.*

Proof. Say $0 \leq k \leq n$, and let $f : S^k \rightarrow |\Delta|$ be any map. By the Cellular Approximation Theorem, there is a map $f' : S^k \rightarrow |\Delta|$ with $f'(S^k) \subset |\Delta^k|$ and with $f \simeq f'$. Let $F : S^k \times I \rightarrow |\Delta|$ be a homotopy from f to f' .

Now since $\Delta^{n+1} = \Gamma^{n+1}$ and $k \leq n$, we can consider f' as a map $S^k \rightarrow |\Gamma|$. Since Γ is n -connected, f' extends to a map $f'' : B^{k+1} \rightarrow |\Gamma|$. Applying the Cellular Approximation Theorem again yields a map $f''' : B^{k+1} \rightarrow |\Gamma^{k+1}| = |\Delta^{k+1}|$. This map is homotopic (rel S^k) to f' , and thus is an extension of f' to B^k . Combining the homotopy F with the map f''' now yields an extension of f to B^{k+1} . $\square$

Note that Lemma 3.1.2 could have been stated, “A simplicial complex is n -connected $\iff$ its $(n+1)$ -skeleton is n -connected.”

We now introduce the concept of an *atomized poset*. Essentially, an atomized poset is a poset in which each set of minimal elements “generates” a well-defined element of the poset (or generates the whole poset).

Definition 3.1.3 *We call a poset P atomized if for each set $S \subset P$ consisting of minimal elements, the subposet $P_{\geq S} = \{p \in P : p \geq s \ \forall s \in S\}$ either is empty or has a minimum element.*

We call the minimal elements of P atoms, and denote the set of atoms of P by $\mathcal{A}(P)$. If $S \subset \mathcal{A}(P)$, we say that S generates the minimum element of $P_{\geq S}$ (or that S generates P if $P_{\geq S}$ is empty), and we write $\langle S \rangle$ for the element generated by S ($\langle S \rangle = P$ is allowed).

It is easy to check that the coset poset and the subgroup poset (the poset of non-trivial, proper subgroups) of a finite group are atomized. The definition of generation coincides with the standard group theoretic definitions (where the coset generated by elements $x_1, \dots, x_n \in G$ is $x_1 \langle x_1^{-1} x_2, \dots, x_1^{-1} x_n \rangle$).

For many purposes, the coset poset is too large to analyze. The following lemma shows that, up to homotopy, we can replace any atomized poset P with a smaller simplicial complex, $\mathcal{M}(P)$. This complex has many fewer vertices, although it has much higher dimension, and will play a crucial role in our analysis of the coset poset.

Lemma 3.1.4 *Let P be an atomized poset and let $\mathcal{M}(P)$ denote the simplicial complex with vertex set $\mathcal{A}(P)$ and with a simplex for each set $S \subset \mathcal{A}(P)$ s.t. $\langle S \rangle \neq P$. Then $\Delta(P) \simeq \mathcal{M}(P)$.*

Proof. Consider the *minimal cover* of P , i.e. the collection of all cones $P_{\geq x}$ with $x \in \mathcal{A}(P)$. We claim that this is a contractible cover of $\Delta(P)$. Indeed, if $S \subset \mathcal{A}(P)$ and $\langle S \rangle \neq P$, then $\cap_{s \in S} P_{\geq s} = P_{\geq \langle S \rangle} \simeq \{*\}$. So each intersection is either empty or contractible, and the Nerve Theorem tells us that $\Delta(P)$ is homotopy equivalent to the nerve of this cover, which is exactly $\mathcal{M}(P)$. $\square$

When $P = \mathcal{C}(G)$ for some finite group G , we denote $\mathcal{M}(\mathcal{C}(G))$ by $\mathcal{M}(G)$. This complex has vertex set G and a simplex for each subset of G contained in a proper coset.

Our main theorem in this section is the following:

Theorem 3.1.5 *Let P be an atomized poset such that no k atoms generate P . Then P is $(k-2)$ -connected.*

Proof. By the previous lemma, it suffices to check that $\mathcal{M}(P)$ is $(k-2)$ -connected. Since no k atoms generate P , any k atoms form a simplex in $\mathcal{M}(P)$. So $\mathcal{M}(P)^{k-1}$ is just the $(k-1)$ -skeleton of the full simplex on the vertex set $\mathcal{A}(P)$. Since any simplex is contractible, Lemma 3.1.2 shows that $\mathcal{M}(P)$ is $(k-2)$ -connected. $\square$

We have the following immediate corollary of Theorem 3.1.6

Corollary 3.1.6 *Let G be a finite group in which any k elements generate a proper subgroup. Then $S(G)$ is $(k-2)$ -connected and $\mathcal{C}(G)$ is $(k-1)$ -connected. In fact, $S(G)$ is $(k-2)$ -connected so long as any k elements of prime order generate a proper subgroup.*

Corollary 3.1.6 specializes to the following result giving conditions under which $\mathcal{C}(G)$ (or any atomized poset) is simply connected.

Corollary 3.1.7 *If G is a finite group in which any two elements generate a proper subgroup, then $\mathcal{C}(G)$ is simply connected. More generally, if P is an atomized poset and no three atoms generate P , then $|P|$ is simply connected.*

As we will see in the next chapter, this theorem does not characterize finite groups with simply connected coset posets. In fact, the first non-abelian simple group, A_5 , affords a counter-example.

We now give an alternate proof of Corollary 3.1.7. This proof will rely solely on the combinatorial structure of $\Delta(P)$ itself (as opposed to that of $\mathcal{M}(P)$). First we introduce some terminology.

By an edge path we mean a map $f : I \rightarrow \Delta(P)^1$ s.t. for some n the restrictions $f|_{[k/n, (k+1)/n]}$ are just a linear shifts of the canonical homeomorphisms $I \rightarrow |\{p_i, p_j\}|$ (where the sets $\{p_i, p_j\}$ are 2-simplices of $\Delta(P)$ and $k = 0, \dots, n-1$). In a simple, closed edge path we require further that $f(0) = f(1)$ and that the p_i are otherwise distinct. From now on we will refer to simple, closed edge paths simply as cycles.


Alternate Proof of Corollary 3.1.7.

Let P be an atomized poset in which no three atoms generate. First, note that $\Delta(P)$ is connected since every element is connected by an edge to an atom, and between any two atoms $a_1, a_2 \in \mathcal{A}(P)$ we have the path $a_1 < \langle a_1, a_2 \rangle > a_2$. We now need to show that $\pi_1(|P|) = 1$, and it will suffice to show that any cycle is null-homotopic (this is a consequence of the Simplicial Approximation Theorem.)

Our first step will be to consider *reduced* cycles. Any cycle can be represented by listing the vertices through which it passes. Let C be a cycle and say C passes through $p_1, p_2, \dots, p_n = p_1$ (in that order). For each i we have either $p_i < p_{i+1}$ or $p_i > p_{i+1}$, and we call a cycle *reduced* if these inequalities alternate direction (they must alternate at the base-point as well).


We claim that any cycle is homotopic to a reduced cycle. If, in a cycle C , some inclusion does not alternate, we have a chain $p_i < p_{i+1} < p_{i+2}$ (or $p_i > p_{i+1} > p_{i+2}$) and removing p_i from the sequence yields a homotopic loop (since this chain forms a 2-simplex). (To be precise, the subscripts above must be taken modulo the length of the cycle; we will continue this abuse of notation throughout the proof.) Repeating this process eventually yields a reduced cycle.

We can simplify any cycle even further. Say $C = (p_1, \dots, p_n)$ is a reduced cycle. We will call a vertex p_i *lower* if $p_i < p_{i+1}$. We call C *atomic* if every lower vertex of C is an atom. Any reduced cycle (and hence any cycle) is homotopic to an atomic cycle, since we may simply replace each lower vertex p_i with an atom $a_i < p_i$ (see the diagram).




Note that both triangles in the above diagram are 2-simplices in $\Delta(P)$.

To complete the proof we must show that any atomic cycle C is null-homotopic. We will proceed by induction on the length of C , which must be even (since C is reduced). The shortest (non-constant) atomic cycle has length 4, and can be written $C = (a_1 < p > a_2 < q > a_1)$. To show that C is null-homotopic, simply consider the point $\langle a_1, a_2 \rangle$. If $\langle a_1, a_2 \rangle \in \{p, q\}$, then it is easy to check that C is null-homotopic, and otherwise the following diagram shows that C is null-homotopic (again, each triangle is a 2-simplex).



Now we may assume that the length of C is greater than 4, so that C contains a path of the form $\rho = (a_1 < p > a_2 < q > a_3)$ (with the a_i atoms). The diagram below shows how to construct a homotopy of ρ with the path $(a_1 < \langle a_1, a_2, a_3 \rangle > a_3)$, assuming that the vertices $a_1, a_2, a_3, p, q, \langle a_1, a_2 \rangle, \langle a_2, a_3 \rangle$ and $\langle a_1, a_2, a_3 \rangle$ are all distinct. (Note that, by hypothesis, $\langle a_1, a_2, a_3 \rangle \neq P$.)



If $\langle a_1, a_2 \rangle = p$ or $\langle a_2, a_3 \rangle = q$, there is no real problem. Otherwise, it is easy to check that if two of these vertices were equal, we would have either $a_1 < q$ or, symmetrically, $a_3 < p$. If $a_1 < q$ then the cycle $(a_1 < p > a_2 < q > a_1)$ is null-homotopic (since it is an atomic cycle of length four) and hence its sub-paths $(a_1 < p > a_2 < q)$ and $(a_1 < q)$ are homotopic. This implies that ρ is homotopic to the path $(a_1 < q > a_3)$.

In any case, we see that ρ is homotopic (relative to its endpoints) to a shorter path, and hence the cycle C is homotopic to a shorter cycle. In addition, this shorter cycle is still atomic, and repeating the process will eventually provide a null-homotopy of C . $\square$

We now present a result guaranteeing simple-connectivity of $\mathcal{C}(G)$ under weaker conditions than those of Corollary 3.1.7. First we need a simple lemma (Brown [7, Proposition 14]).

Lemma 3.1.8 *$\mathcal{C}(G)$ is connected unless G is cyclic of prime-power order.*

Proof. If G is non-cyclic, then any proper coset xH connects to the identity via the path $xH \geq \{x\} \leq \langle x \rangle \geq \{1\}$. Now say G is cyclic but not of prime power order. If x does not generate G , then any coset xH connects to the identity as before. If x is a generator, then we may write $x = yz$ where y

and z are not generators (this is possible because G decomposes as a direct product in a non-trivial manner). Then any coset xH connects to $\{1\}$ via the path

$$xH \geq \{x = yz\} \leq y\langle z \rangle \geq \{y\} \leq \langle y \rangle \geq \{1\}.$$

If G is cyclic of prime-power order, then G has a unique maximal subgroup M , and $\mathcal{C}(G)$ is the disjoint union of the cones $\mathcal{C}(G)_{\leq xM}$. If $G \cong \mathbb{Z}/p^n$, then $(G : M) = p$ and we see that $\mathcal{C}(G)$ is homotopy equivalent to a discrete space with p points. $\square$

Theorem 3.1.9 *Let G be a finite, non-cyclic group in which the following condition holds: For any $x, y \in G$ s.t. $\langle x, y \rangle = G$ there exists $z \in G$ s.t. $\langle z, x \rangle \neq G$, $\langle z, y \rangle \neq G$ and $\langle zx, zy \rangle \neq G$. Then $\mathcal{C}(G)$ is simply connected.*

Proof. Let G be a group satisfying the hypotheses of the theorem. We will show that $\pi_1(\mathcal{M}(G)) = 0$. First, consider the 1-skeleton of $\mathcal{M}(G)$. Since G is non-cyclic, any two elements x, y are contained in a proper coset, namely $x\langle x^{-1}y \rangle$. Thus in $\mathcal{M}(G)$ all possible edges between vertices exist.

We wish to apply Theorem 2.1.9 to $\mathcal{M}(G)$, so we must choose a maximal tree in $\mathcal{M}(G)$ ¹. Since all possible edges exist, we can choose our maximal tree T to be the star at the vertex 1, i.e. T consists of all edges of the form $\{1, x\}$ with $x \in G$. In order to show that $\pi_1(\mathcal{M}(G))$ is trivial, we will use the relations given in Theorem 2.1.9 to show that each generator is trivial.

There are two cases. First, if x and y do not generate G , then the subgroup $\langle x, y \rangle$ is a coset containing 1, x and y , and thus these vertices form a simplex in $\mathcal{M}(G)$. So we obtain the relation $(x, y)(y, 1)(1, x) = 1$, which forces (x, y) to be trivial.

Next, consider the case in which $\langle x, y \rangle = G$. Let z be the element guaranteed by the hypotheses. Then, as above, the generators corresponding to the edges $\{x, z^{-1}\}$ and $\{y, z^{-1}\}$ are trivial (note that $\langle x, z \rangle \neq G \iff \langle x, z^{-1} \rangle \neq G$). Furthermore, since $\langle zx, zy \rangle \neq G$ there is a simplex $\{x, y, z^{-1}\}$ in $\mathcal{M}(G)$, and we have the relation $(x, y)(y, z^{-1})(z^{-1}, x) = 1$, which again forces (x, y) to be trivial.

Thus all generators for $\pi_1(\mathcal{M}(G))$ are trivial, and hence $\mathcal{M}(G)$ and $\mathcal{C}(G)$ are simply connected. $\square$

In the next chapter we will apply this result to show that $\mathcal{C}(A_5)$ is simply connected.

3.2 Mayer-Vietoris Sequences and $H_*(\mathcal{C}(G))$

We will now present some results based on Mayer-Vietoris sequences for the homology of $\mathcal{C}(G)$.

From here on, we let $H_n(X)$ denote the homology of X with coefficients in $\mathbb{Z}$, and we let $\tilde{H}_n(X)$ denote reduced homology (again with coefficients in $\mathbb{Z}$).

Theorem 3.2.1 *Let G be a finite non-cyclic group with a cyclic maximal subgroup M of prime-power order. Let $o(M) = p^n$. Then $H_1(\mathcal{C}(G))$ has rank at least $(p-1)(G:M)$, and in particular $\mathcal{C}(G)$ is not simply connected.*

Proof. Let $\{x_i\}_{i \in I}$ be a set of (left) coset representatives for M . Let $X = \Delta(\mathcal{C}(G))$, $Y = \Delta(\mathcal{C}(G) - \{x_i M\}_{i \in I})$, and $Z = \Delta(\cup_{i \in I} \mathcal{C}(G)_{\leq x_i M})$. Then we have $X = Y \cup Z$, since any chain in $\mathcal{C}(G)$ either contains $x_i M$ for some $i \in I$, and lies in Z , or does not contain any coset of M , and lies in Y . Thus we have a Mayer-Vietoris sequence

$$\cdots \longrightarrow \tilde{H}_1(X) \xrightarrow{f} \tilde{H}_0(Y \cap Z) \xrightarrow{h} \tilde{H}_0(Y) \oplus \tilde{H}_0(Z) \longrightarrow \tilde{H}_0(X)$$

on (reduced) homology groups.

Note that X is connected because G is non-cyclic (Lemma 3.1.8), and hence $\tilde{H}_0(X) = 0$. Also, Z is a union of $(G:M)$ cones, one for each $i \in I$, and hence $\tilde{H}_0(Z) \cong \mathbb{Z}^{(G:M)-1}$. In addition, we claim that Y is connected so that $\tilde{H}_0(Y) = 0$. Substituting these values into the above sequence yields an exact sequence

$$\cdots \longrightarrow \tilde{H}_1(X) \xrightarrow{f} \tilde{H}_0(Y \cap Z) \xrightarrow{h} \mathbb{Z}^{(G:M)-1} \longrightarrow 0,$$

and thus $h : \tilde{H}_0(Y \cap Z) \longrightarrow \mathbb{Z}^{(G:M)-1}$ is a surjection. Now,

$$Y \cap Z = \Delta(\cup_{i \in I} \mathcal{C}(G)_{< x_i M}) \cong \coprod_{i \in I} \mathcal{C}(M).$$

As discussed in the proof of Lemma 3.1.8, the coset poset of $M \cong \mathbb{Z}/p^n$ is disconnected, and in fact it has p contractible components (the cones under the cosets of the unique maximal subgroup). Thus $\tilde{H}_0(Y \cap Z) \cong \mathbb{Z}^{p(G:M)-1}$, and thus $\text{Im } f = \text{Ker } h \cong \mathbb{Z}^{p(G:M)-1 - ((G:M)-1)} = \mathbb{Z}^{(p-1)(G:M)}$ and we see that the rank of $H_1(X)$ is at least $(p-1)(G:M)$, as desired.

To complete the proof we must show that Y is connected. Consider a coset xH , where $H \neq M$. If $\langle x \rangle \neq M$, then we have a path

$$xH \geq \{x\} \leq \langle x \rangle \geq \{1\}$$

connecting xH to the identity (note that we have assumed G is non-cyclic, so that $\langle x \rangle \in Y$). Next, if $\langle x \rangle = M$ then choose some $g \in G$, $g \notin M$. We now have a path

$$xH \geq \{x\} \leq x\langle g \rangle \geq \{xg\} \leq \langle xg \rangle \geq \{1\}$$

connecting xH to the identity (note that $\langle xg \rangle \neq M$, since if $xg \in M$ then $g \in x^{-1}M = M$, contrary to our choice of g). $\square$

The above result does not extend to cyclic groups, as we will see in a moment. The basic problem is that the subcomplex Y is no longer connected. Nevertheless, the coset poset of a cyclic group with maximal subgroup isomorphic to $\mathbb{Z}/p^n$ is still not simply connected. Any group of this type may be written as $\mathbb{Z}/q \times \mathbb{Z}/p^n$, with q prime, and now [7, Lemma 5] implies that $\mathcal{C}(G) \cong |\mathcal{C}(\mathbb{Z}/q)| * |\mathcal{C}(\mathbb{Z}/p^n)|$ (where $*$ denotes the join). The latter is (homotopy equivalent to) a bipartite graph with $q + p$ vertices and pq edges. Modding out a maximal tree in this graph and applying Van Kampen's Theorem shows that $H_1(\mathcal{C}(G)) \cong \mathbb{Z}^{(p-1)(q-1)}$. (For G to satisfy the conclusion of the theorem, we would need the rank of H_1 to be at least $(p-1)q$.)

Question 3.2.2 *Does Theorem 3.2.1 characterize (non-cyclic) finite groups with simply connected coset posets?*

We expect the answer to be no because, as we will show later in this section, Theorem 3.2.1 applies only to a small class of solvable groups. In addition, Corollary 3.2.4 may allow one to find other finite groups with non-simply connected coset posets.

The following theorem shows a (somewhat weak) connection between the coset poset and the subgroup poset of a finite group. In Chapter 5, we will discuss other possible connections.

Theorem 3.2.3 *Let G be a finite group, and for $g \in G$, let $\mathcal{C}(G)_g$ denote the poset $\mathcal{C}(G) - \{g\}$.*

If there exists $g \in G$ with $\tilde{H}_n(\mathcal{C}(G)_g) = 0$, then there is a surjection $\tilde{H}_{n+1}(\mathcal{C}(G)) \rightarrow \tilde{H}_n(S(G))$.

Note that for any $g, h \in G$, the posets $\mathcal{C}(G)_g$ and $\mathcal{C}(G)_h$ are isomorphic (consider the map given by left-multiplication by hg^{-1}). In particular, $\mathcal{C}(G)_g \cong \mathcal{C}(G)_1 \cong S(G)$.

Proof. Let $X = \Delta(\mathcal{C}(G))$, $Y = \Delta(\mathcal{C}(G)_{\geq \{g\}})$, and $Z = \Delta(\mathcal{C}(G)_g)$. Note that Y is contractible (so $\tilde{H}_n(Y) = 0$ for all n) and that $Y \cap Z \cong S(G)$.

Since $X = Y \cup Z$, we get a Mayer-Vietoris sequence

$$\cdots \longrightarrow \tilde{H}_{n+1}(X) \xrightarrow{f} \tilde{H}_n(Y \cap Z) \longrightarrow \tilde{H}_n(Y) \oplus \tilde{H}_n(Z) \longrightarrow H_n(X) \longrightarrow \cdots$$

and since $\tilde{H}_n(Y) = \tilde{H}_n(Z) = 0$, the map

$$f : H_{n+1}(X) \rightarrow H_n(Y \cap Z) \cong H_n(S(G))$$

is a surjection. □

Corollary 3.2.4 *Let G be a finite group and let n be the number of path-components of $S(G)$. Then $H_1(\mathcal{C}(G))$ has rank at least $n - 1$. In particular, if $S(G)$ is disconnected then $\mathcal{C}(G)$ is not simply connected.*

Proof. First, if G is cyclic of prime-power order then $\mathcal{C}(G)$ is homotopy equivalent to a discrete set of p points, and hence $H_1(\mathcal{C}(G)) = 0$. Also, G has a unique maximal subgroup in this case, and hence $S(G)$ has one path component.

Now we turn to the case in which G is not cyclic of prime-power order. In light of Theorem 3.2.3 it will suffice to prove that $\mathcal{C}(G)_1$ is connected. First, consider the case in which G is not cyclic. Choose some element $x \in G$, $x \neq 1$. Then any vertex yH in Y can be connected to $\{x\}$ via the path

$$yH \geq \{y\} \leq x\langle x^{-1}y \rangle \geq \{x\}$$

(note that G is non-cyclic so $x\langle x^{-1}y \rangle \in Y$, and $yH \neq \{1\}$ so we may assume $y \neq 1$).

If G is cyclic (but not of prime-power order), let x be a generator for G . Then any coset in G can be written as $x^n\langle x^m \rangle$, for some $n, m > 0$. Also, note that since $o(G)$ has at least two prime factors, G decomposes as a (non-trivial) direct product and hence any element in G can be written as a product of two non-generators. Specifically, we may choose k, l s.t. $x^{n-1} = x^k x^l$ and

$\langle x^k \rangle, \langle x^l \rangle$ are proper subgroups of G . Using this decomposition, we have the following path joining $x^n \langle x^m \rangle$ to $\{x\}$:

$$x^n \langle x^m \rangle \geq \{x^n = x^{k+1} x^l\} \leq x^{k+1} \langle x^l \rangle \geq \{x^{k+1}\} \leq x \langle x^k \rangle \geq \{x\}.$$

Thus Y is connected and the proof is complete. $\square$

Corollary 3.2.4 does not characterize finite groups with simply connected coset posets. The quaternion group Q_8 provides a counter-example, since $S(Q_8)$ is contractible (it is a cone on $\langle -1 \rangle$) and yet $\pi_1(\mathcal{C}(G)(Q_8)) \cong \mathbb{Z} * \mathbb{Z} * \mathbb{Z}$, as will be shown in the next chapter (Section 4.1).

Question 3.2.5 *For which finite groups G is $S(G)$ disconnected? In particular, do there exist non-solvable groups with disconnected subgroup lattices?*

We now turn to the question of which groups satisfy the conditions of Theorem 3.2.1. In the case where G is a p -group, these groups have been classified (see Brown [5, Theorem 4.1, p. 98]).

We now turn to the case in which $o(G)$ has at least two prime divisors, p and q . First, the following result can be found in Herstein [13]:

Theorem 3.2.6 *Let G be a finite group with an abelian maximal subgroup. Then G is solvable.*

In our case, something stronger may be shown. Our proof of the following result is based on Herstein's proof of Theorem 3.2.6.

Theorem 3.2.7 *Let G be a finite group and assume p and q are distinct primes dividing $o(G)$. Assume further that G has a maximal subgroup M which is an abelian p -group. Then either $G \cong M \rtimes \mathbb{Z}/q$ or $G \cong Q \rtimes M$, where Q is the (unique) q -Sylow subgroup of G .*

Note that solvability of G follows immediately from this theorem, since Q and M are each nilpotent.

In proving the theorem, we will need a result due to Burnside (see [19, p. 289]) and a lemma from Herstein [13]. Herstein's proof is elementary, and will be included for completeness.

Theorem 3.2.8 (Burnside's Theorem) *Let G be a finite group and let P be a p -Sylow subgroup of G which lies in the center of its normalizer. Then G is p -nilpotent, that is there exists a subgroup $T \triangleleft G$ s.t. $T \cap P = \{1\}$ and $TP = G$ (hence $G \cong T \rtimes P$).*

Lemma 3.2.9 *Let $\mathcal{A}$ be an abelian group of automorphisms of a finite group G . Assume that each non-identity element $\alpha \in \mathcal{A}$ fixes only the identity in G . Then for each prime p dividing $o(G)$, there exists a unique Sylow p -subgroup S_p of G which is invariant under the action of $\mathcal{A}$.*

Proof. Let $\alpha \in \mathcal{A}$ be a non-trivial automorphism of G . Consider the set $\{g^{-1}\alpha(g) : g \in G\}$. We claim that this set is all of G . Indeed, if $g^{-1}\alpha(g) = h^{-1}\alpha(h)$ then $hg^{-1} = \alpha(hg^{-1})$ and thus hg^{-1} is fixed by α . By hypothesis we now have $hg^{-1} = 1$, and thus $h = g$.

Now, fix a prime p dividing $o(G)$ and a Sylow p -subgroup P . Then $\alpha(P) = xPx^{-1}$ for some $x \in G$, and (by the discussion above) there exists $y \in G$ s.t. $y^{-1}\alpha(y) = x^{-1}$. So we have

$$\alpha(yPy^{-1}) = \alpha(y)\alpha(P)\alpha(y^{-1}) = \alpha(y)xPx^{-1}\alpha(y^{-1}) = yPy^{-1}.$$

Thus yPy^{-1} is a Sylow p -subgroup fixed by α . We claim that in fact, this is the only Sylow p -subgroup fixed by α . Assuming uniqueness, we now complete the proof. Letting $S_p = yPy^{-1}$, we see that for any $\beta \in \mathcal{A}$

$$\alpha\beta(S_p) = \beta\alpha(S_p) = \beta(S_p).$$

(Recall that $\mathcal{A}$ is abelian.) Thus $\beta(S_p)$ is also fixed by α and must be S_p . So S_p is fixed by all of $\mathcal{A}$.

We will now prove uniqueness of S_p . If S'_p is another p -Sylow subgroup fixed by α , we have $gS_pg^{-1} = S'_p$ for some $g \in G$, and $g^{-1}\alpha(g) \in N(S_p)$ because

$$g^{-1}\alpha(g)S_p\alpha(g^{-1})g = g^{-1}\alpha(gS_pg^{-1})g = g^{-1}\alpha(S'_p)g = g^{-1}S'_pg = S_p.$$

Now, note that α induces an automorphism of $N(S_p)$ which fixes only the identity, and thus all elements of $N(S_p)$ have the form $n^{-1}\alpha(n)$ for some $n \in N(S_p)$. Since $g^{-1}\alpha(g) \in N(S_p)$, this means $g^{-1}\alpha(g) = n^{-1}\alpha(n)$ for some $n \in N(S_p)$, and hence $g = n$ (as was shown at the start of the proof). So $g \in N(S_p)$, which means $S'_p = S_p$. This shows that S_p is the unique Sylow p -subgroup fixed by α , completing the proof. $\square$

Proof of Theorem 3.2.7. Let G be a finite group satisfying the hypotheses of the theorem. If $M \triangleleft G$, then G/M has no non-trivial subgroups, and hence $G/M \cong \mathbb{Z}/q$. Thus $o(G) = q o(M)$, and hence the q -Sylow subgroups of G are isomorphic to $\mathbb{Z}/q$ and we have $G \cong M \rtimes \mathbb{Z}/q$.

Next, say $M \ntriangleleft G$. We will proceed by induction on $o(G)$. Since $p, q | o(G)$, the base case occurs when $o(G) = pq$. In this case $G \cong \mathbb{Z}/q \rtimes \mathbb{Z}/p$, since $M \ntriangleleft G$ (this follows from an easy argument using the Sylow theorems.) We now assume that for any group H with $o(H) < o(G)$, $p, q | o(H)$ and with an abelian maximal subgroup P which is a p -group, $P \ntriangleleft H$ implies $H \cong Q \rtimes P$ (where Q is the Sylow q -subgroup of H).

Now, since M is maximal, $N(M) = M$ and since M is abelian Burnside's Theorem applies. So there is a subgroup $T \triangleleft G$ with $G \cong T \rtimes M$, and it remains to show that T is a q -group.

We will consider two cases. First we assume that for any $g \in G$, $g \notin M$, $(g^{-1}Mg) \cap M = \{1\}$. Consider the action of M on T by conjugation. We claim that for any $m \in M$ ($m \neq 1$), the only element of T fixed under conjugation by m is 1. Indeed, if $t \in T$ and $m^{-1}tm = t$, then $tmt^{-1} = m \in M \cap (tMt^{-1})$. Since $m \neq 1$, $M \cap (tMt^{-1}) \neq \{1\}$ and hence $t \in M$, which implies $t = 1$ (since $t \in T$ and $T \cap M = \{1\}$). Thus we may apply Herstein's lemma to the (abelian) group of automorphisms of T induced by M . This gives us a unique Sylow q -subgroup Q of T normalized by M (note that $q | o(T)$ because $q | o(G)$ and $G \cong T \rtimes M$). Now we have $Q, M \subset N(Q)$, so $Q \triangleleft G$ by maximality of M . Hence QM is a subgroup of G , and since M is maximal we have $QM = G$. This shows that $G \cong Q \rtimes M$.

Finally we must consider the case in which $M \triangleleft G$ and there exists $g \in G$, $g \notin M$, with $(g^{-1}Mg) \cap M \neq \{1\}$. Let $W = (g^{-1}Mg) \cap M$, and choose $w \in W$, $w \neq 1$. Note that $W < M$, since $N(M) = M$ and $g \notin M$. We now have $\langle w \rangle \triangleleft M, g^{-1}Mg$ (since these groups are abelian), which implies $\langle w \rangle \triangleleft G$ (by maximality of M).

Consider $G/\langle w \rangle$. Since $\langle w \rangle \leq W < M$, $M/\langle w \rangle$ is an abelian p -group, and is maximal in $G/\langle w \rangle$. Also, we have $p, q | o(G/\langle w \rangle)$, and since $M \triangleleft G$, $M/\langle w \rangle \triangleleft G/\langle w \rangle$. So $G/\langle w \rangle$ is a group of smaller order satisfying the conditions of the induction hypothesis, and we may assume that $G/\langle w \rangle \cong \bar{Q} \rtimes M/\langle w \rangle$, where $\bar{Q}$ is the Sylow q -subgroup of $G/\langle w \rangle$. Notice that p and q are the only prime factors of $o(G/\langle w \rangle)$, and since $\langle w \rangle$ is a p -group, these must be the only prime factors of $o(G)$ as well. Finally, since M is maximal it must be a Sylow p -subgroup of G , and hence T is a q -group, as desired. $\square$

Corollary 3.2.10 *Let G be a group with maximal subgroup $M \cong \mathbb{Z}/p^n$, for some prime p and some $n > 0$. If G is not a p -group, then there is a prime $q \neq p$ such that either $G \cong \mathbb{Z}/p^n \rtimes \mathbb{Z}/q$ or $G \cong Q \rtimes \mathbb{Z}/p^n$ (with Q a q -group).*

3.3 Applications to Group Theory

We now present group theoretic applications of the results in the previous section. First we will need to discuss a result of Brown [7, Proposition 11], describing the homotopy type of the coset poset of a solvable group.

Definition 3.3.1 *Let G be a finite group, and let P be the poset of normal subgroups of G (including $\{1\}$ and G), ordered by inclusion. The maximal chains in P are called chief series for G .*

If $\{1\} = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_k = G$ is a chief series for G , we call an element G_i , $i = 1, \dots, k$, complemented if G_i/G_{i-1} has a complement in G/G_{i-1} .

Theorem 3.3.2 *Let G be a finite solvable group, and let*

$$\{1\} = G_0 \triangleleft G_1 \triangleleft \cdots \triangleleft G_k = G$$

be a chief series for G . Then $\mathcal{C}(G)$ is homotopy equivalent to a bouquet of spheres, each of dimension $d-1$ where d is the number of indices i , $1 \leq i \leq k$, s.t. G_i is complemented in the above chief series.

Corollary 3.3.3 *Let G be a finite group with maximal subgroup $M \cong \mathbb{Z}/p^n$, for some prime p and some $n > 0$. Then in any chief series for G , exactly one proper subgroup is complemented.*

Proof. The results of the previous section imply that G is solvable, so Theorem 3.3.2 implies that $\mathcal{C}(G)$ has the homotopy type of a bouquet of spheres. Since $\mathcal{C}(G)$ is not simply connected (by Theorem 3.2.1) the dimension of these spheres must be 1. So $d = 2$, and there is exactly one complemented factor (other than G itself) in any chief series for G . $\square$

Similarly, we have:

Corollary 3.3.4 *Let G be a solvable finite group with disconnected subgroup poset. Then in any chief series for G , exactly one proper subgroup is complemented.*

3.4 Group Extensions

In this section we will consider the coset poset of a group G which may be written as an extension

$$N \hookrightarrow G \longrightarrow G/N.$$

First, we characterize (non-trivial) direct products with simply connected coset posets (Corollary 3.4.3). The proof employs a result from Brown [7, Lemma 5] on direct products, which we generalize to semi-direct products (Theorem 3.4.4). Lastly, we employ another result from [7] to show that if a group G has a quotient G/N whose coset poset is simply connected, then $\mathcal{C}(G)$ is simply connected as well. These results show, in particular, that there are infinitely many non-solvable groups with simply-connected coset posets.

Definition 3.4.1 *We call a coset $C \in \mathcal{C}(G \times H)$ saturating if it surjects onto each factor under the natural projections from $G \times H$ to G and H . The poset of non-saturating cosets of $G \times H$ will be denoted $\mathcal{C}_0(G \times H)$.*

Theorem 3.4.2 (Brown) *For any finite groups G and H ,*

$$\mathcal{C}_0(G \times H) \simeq \mathcal{C}(G) * \mathcal{C}(H),$$

where $$ denotes the join operation.*

For the purposes of this section, we simply regard $P * Q$ as the topological space $|P| * |Q|$ (for any posets P and Q). For an alternate (and compatible) viewpoint, see Quillen [18].

Corollary 3.4.3 *Let G and H be non-trivial finite groups. Then $\mathcal{C}(G \times H)$ is simply connected if and only if at least one of these groups is not cyclic of prime-power order.*

Proof. If both groups are cyclic of prime-power order, it is easy to check that there are just two split factors in any chief series for $G \times H$ (in the sense of Theorem 3.3.2) and the desired result then follows from that theorem.

In the other direction, assume WLOG that G is not cyclic of prime-power order (so $\mathcal{C}(G)$ is connected). Then $\mathcal{C}_0(G \times H)$ is simply connected:

by Theorem 3.4.2, this space is (up to homotopy) the join of $\mathcal{C}(H)$ with the connected space $\mathcal{C}(G)$, and the join of a non-empty space with a connected space is simply connected (see Milnor [16]).

Now we consider the effect of adding saturating cosets to $\mathcal{C}_0(G \times H)$. Consider a minimal element xK of the poset $\mathcal{C}(G \times H) - \mathcal{C}_0(G \times H)$. Adding this to $\mathcal{C}_0(G \times H)$ has the effect of coning off a copy of $\mathcal{C}(K) \cong \mathcal{C}(G)_{\leq xK}$, which we claim is connected. Connectivity of $\mathcal{C}(K)$ results from the fact that K surjects onto G and H : since one of these is not cyclic of prime-power order, neither is K and hence $\mathcal{C}(K)$ is connected (by Lemma 3.1.8). Now, coning off a connected subset of a simply connected space yields another simply connected space (this follows from Van Kampen's Theorem, or can be proven directly).

We may now continue in this manner, adding in minimal elements from the poset of remaining cosets, and at each stage we simply cone off a copy of $\mathcal{C}(K)$, where $K \leq G \times H$ surjects onto G and onto H . Thus we see that $\mathcal{C}(G \times H)$ is simply connected. $\square$

We will now generalize Theorem 3.4.2 to the case of a semi-direct product.

For any semi-direct product $G \rtimes H$ (with G and H finite), let $f : G \rtimes H \rightarrow G$ be the function $f(g, h) = g$, and let $\pi : G \rtimes H \rightarrow H$ be the quotient map. We call a coset $xT \in \mathcal{C}(G \rtimes H)$ *saturating* if $\pi(xT) = H$ and the only H -invariant subgroup of G that contains $f(T)$ is G itself.

Theorem 3.4.4 *Let G and H be finite groups and consider a semi-direct product $G \rtimes H$. Let $\mathcal{C}_0(G \rtimes H)$ be the poset of all non-saturating cosets and let $\mathcal{C}_H(G)$ denote the poset of all cosets of proper, H -invariant subgroups of G . Then $\mathcal{C}_0(G \rtimes H)$ is homotopy equivalent to the join $\mathcal{C}_H(G) * \mathcal{C}(H)$.*

Proof. Let $\mathcal{C}^+(H)$ denote the set of all cosets in H (including H itself) and let $\mathcal{C}_H^+(G) = \mathcal{C}_H(G) \cup \{G\}$. Then if $\mathcal{C}_{00}(G \rtimes H)$ denotes the set of all proper cosets of the form $(g, h)I \rtimes K$ (with $I \leq G$ invariant and $K \leq H$), one checks that the map

$$(x, y)I \rtimes K \mapsto (xI, yK)$$

is a well-defined poset isomorphism

$$\mathcal{C}_{00}(G \rtimes H) \xrightarrow{\cong} \mathcal{C}_H^+(G) \times \mathcal{C}^+(H) - \{(G, H)\}.$$

The latter has geometric realization homotopy equivalent to $\mathcal{C}_H(G) * \mathcal{C}(H)$ (see Quillen [18, Proposition 1.9]). Finally, we have an increasing poset map

Φ from $\mathcal{C}_0(G \rtimes H)$ onto $\mathcal{C}_{00}(G \rtimes H)$ given by

$$\Phi((x, y)T) = (x, y)\hat{f}(T) \rtimes \pi(T)$$

where $\hat{f}(T)$ is the smallest invariant subgroup containing $f(T)$ (i.e. $\hat{f}(T)$ is the intersection of all invariant subgroups containing $f(T)$). Corollary 2.2.13 shows that this map is a homotopy equivalence between $\mathcal{C}_0(G \rtimes H)$ and $\Phi(\mathcal{C}_0(G \rtimes H)) = \mathcal{C}_{00}(G \rtimes H)$ (with homotopy inverse the inclusion map $i : \mathcal{C}_{00}(G \rtimes H) \rightarrow \mathcal{C}_0(G \rtimes H)$). $\square$

With this theorem in hand, one ought to be able to show that many semi-direct products have simply connected coset posets. In light of the results of Section 3.2, the collection will not be quite the same as in the case of direct products, but hopefully it will be relatively large.

We now turn to the case of a general group extension, and prove a simple corollary to the following result of Brown [7, Proposition 10].

Theorem 3.4.5 *For any finite group G and any normal subgroup $N \triangleleft G$ there is a homotopy equivalence*

$$\mathcal{C}(G) \longrightarrow \mathcal{C}(G/N) * \mathcal{C}(G, N),$$

where $\mathcal{C}(G, N)$ denotes the poset of saturating cosets, i.e. cosets $C \in \mathcal{C}(G)$ s.t. $q(C) = G/N$.

Corollary 3.4.6 *Let G be a finite group with quotient H , and assume that $\mathcal{C}(H)$ is simply connected. Then $\mathcal{C}(G)$ is simply connected as well.*

Proof. Choose $N \triangleleft G$ s.t. $G/N \cong H$. Then Theorem 3.4.5 tells us that $\mathcal{C}(G) \simeq \mathcal{C}(H) * \mathcal{C}(G, N)$, and if $\mathcal{C}(H)$ is simply connected then its join with any space is simply connected. $\square$

Presumably, it would be easy to extend the results of this section to deal with higher connectivity, and it would be interesting to explore this further.

Chapter 4

Examples

In this chapter, we will examine the homotopy type of $\mathcal{C}(G)$ for some specific finite groups G . Along the way, we will introduce new techniques for dealing with the coset poset.

The main goal of this chapter is to study the coset posets of two finite simple groups, A_5 and $PSL_2(\mathbb{F}_7)$. Brown's theorem (Theorem 3.3.2), gives a complete description of the homotopy type of the coset poset of any finite solvable group, and but little is known about non-solvable groups. Thus it makes sense to study simple groups, at the opposite end of the spectrum. In particular, we will give two simple proofs of an unpublished result of Shareshian (see [7, p. 1009]) describing the homotopy type of $\mathcal{C}(A_5)$, and we will show that the coset poset of $PSL_2(\mathbb{F}_7)$ is simply connected.

Before moving on to these more complicated results, we present another description of the coset poset and use this description to calculate the homotopy type of $\mathcal{C}(Q_8)$, where Q_8 is the quaternion group. This will show that Corollary 3.2.4 does not characterize groups with simply connected coset posets.

4.1 The Quaternion Group

Let Q_8 denote the standard quaternion group consisting of the 8 elements $\pm 1, \pm i, \pm j, \pm k$. The proper, non-trivial subgroups of Q_8 are just $\langle -1 \rangle = \{\pm 1\}$, $\langle i \rangle = \{\pm 1, \pm i\}$, $\langle j \rangle = \{\pm 1, \pm j\}$, and $\langle k \rangle = \{\pm 1, \pm k\}$. Thus $S(Q_8)$ is

a cone with minimum element $\langle -1 \rangle$, and hence this poset is contractible (so Corollary 3.2.4 does not apply to it).

We will now calculate the homotopy type of $\mathcal{C}(Q_8)$, and show that this poset is not simply connected. To do this, we use the Cross-Cut Theorem [3, Theorem 10.8].

Definition 4.1.1 *Let P be a poset, and let $C \subset P$ be a subposet satisfying:*

1. *C is an antichain (no two elements of C are comparable),*
2. *If σ is a chain in P then there is some $c \in C$ which is comparable to all elements of σ ,*
3. *If $A \subset C$ has an upper (lower) bound in P , then A has a meet (join) in P .*

We call C a cross-cut, and we define $\Gamma(P, C)$, the cross-cut complex associated to C , to be the simplicial complex with vertex set C and with a simplex for each subset of C which is bounded in P .

Lemma 4.1.2 (Cross-Cut Theorem) *Let P be a poset and let $C \subset P$ be a cross-cut. Then $\Delta(P) \simeq \Gamma(P, C)$.*

Proof. The proof is an application of the Nerve Theorem. Consider the subposets $D_x = P_{\geq x} \cup P_{\leq x}$ for $x \in P$. We claim that the subcomplexes $\Delta(D_c)$, $c \in C$, give a contractible cover of $\Delta(P)$.

Condition 2 gives $\Delta(P) = \bigcup_{c \in C} \Delta(D_c)$. Next, say $C' \subset C$. Then C' is an antichain, so if $x \in D_{c'}$ for each $c' \in C'$, we either have $x > c'$ for all $c' \in C'$ or $x < c'$ for all $c' \in C'$. Thus if this intersection is non-empty, C' is bounded in P and hence has either a meet or a join (by condition 3). In either case, this meet or join is comparable to each element in the intersection. So $\bigcap_{c' \in C'} D_{c'}$ is either empty or a cone, hence contractible.

So we have a contractible cover of P , and in fact we see that $\Gamma(P, C)$ is exactly the nerve of this cover. This completes the proof. $\square$

Corollary 4.1.3 *Let G be a finite group and denote by $Pr(G)$ the simplicial complex with vertex set $V = \{xH \in \mathcal{C}(G) : o(H) \text{ is prime}\}$ and with a simplex for each subset of V bounded in $\mathcal{C}(G)$. Then $\mathcal{C}(G) \simeq Pr(G)$.*

Proof. It suffices to show that V is a cross-cut in $\mathcal{C}(G)$ (since $Pr(G)$ is then the cross-cut complex of V). Certainly, V is an antichain since no two subgroups of prime order are comparable. Next, any chain $xH_1 < xH_2 < \cdots < xH_n$ with $H_1 \neq \{1\}$ lies above xP , where P is a subgroup of prime order in H_1 . If $H_1 = \{1\}$, then letting P be a prime-order subgroup of H_2 , we see that xH_i is comparable to xP for each i . Finally, the third condition of a cross-cut is satisfied trivially, since any subset in $\mathcal{C}(G)$ with a lower (upper) bound has a meet (join). $\square$

Claim 4.1.4 $\mathcal{C}(Q_8)$ has the homotopy type of a wedge of three circles, and hence $\pi_1(\mathcal{C}(Q_8)) \cong \mathbb{Z} * \mathbb{Z} * \mathbb{Z}$.

Proof. We will apply Corollary 4.1.3 to Q_8 . Note that the only subgroup of prime order in Q_8 is $\langle -1 \rangle$. The cosets of $\langle -1 \rangle$ are $\{1, -1\}$, $\{i, -i\}$, $\{j, -j\}$ and $\{k, -k\}$, and it is easy to check that any two are contained in a proper coset. So $Pr(Q_8)$ is the complete graph on 4 vertices, and has six edges. A maximal tree in $Pr(Q_8)$ has three edges, and after modding out such a tree we are left with a wedge of three circles. The result now follows from Van Kampen's Theorem. $\square$

4.2 The Coset Poset of A_5

In this section we will consider the smallest example of a group whose coset poset is not determined (up to homotopy) by Brown's theorem (Theorem 3.3.2). This group is A_5 , the first non-solvable group (as well as the first non-abelian simple group). Our goal will be to establish the following theorem.

Theorem 4.2.1 *The coset poset of A_5 has the homotopy type of a bouquet of 1560 two-dimensional spheres.*

It can be checked that there are 1018 proper cosets in A_5 , and hence $\mathcal{C}(A_5)$ has 1018 vertices. Evidently, $\mathcal{C}(A_5)$ is far too large to admit direct analysis.

The proof of Theorem 4.2.1 will proceed in stages. First we will show that $\mathcal{C} = \mathcal{C}(A_5)$ has the homotopy-type of a two-dimensional complex. For this portion of the proof we will work directly with $\mathcal{C}$. Then we will show that $\mathcal{C}$ is simply connected by examining $\mathcal{M}(A_5)$. We will give two proofs

of simple connectivity: the first will be an application of Theorem 3.1.9 and the second will use 2-transitive actions of A_5 . To show that $\mathcal{C}(A_5)$ has the homotopy type of a bouquet of 2-spheres, we appeal to the general result that a k -dimensional complex which is $(k - 1)$ -connected is homotopy equivalent to a bouquet of k -spheres.

The number of spheres in the bouquet can be calculated from the Euler characteristic $\tilde{\chi}(\mathcal{C}(A_5))$. Brown [7] has shown that $\tilde{\chi}(\mathcal{C}(A_5)) = 1560$, and viewing the Euler characteristic as an alternating sum of homology ranks, we see that the number of 2-spheres is exactly 1560.

We will now examine the subgroup lattice of A_5 , in order to describe the chains in $\mathcal{C}$ of length greater than three (i.e. the simplices in $\Delta(\mathcal{C})$ of dimension greater than two). We will consider A_5 together with its standard action on the set $\{1, 2, 3, 4, 5\}$, and we say that a subgroup $H \leq A_5$ has a fixed point if some element of $\{1, 2, 3, 4, 5\}$ is fixed by all elements of H .

We use the notation D_{2n} for the dihedral group of order $2n$ (so $D_{2n} = \mathbb{Z}/n \rtimes \mathbb{Z}/2$).

Claim 4.2.2 *If M is a maximal subgroup of A_5 , then M is isomorphic to one of D_{10} , A_4 , S_3 . There are exactly five copies of A_4 corresponding to the five subsets of $\{1, 2, 3, 4, 5\}$ of cardinality four; there are six copies of D_{10} , each normalizing a different copy of $\mathbb{Z}/5$; and there are ten copies of S_3 , all conjugate to one another in S_5 .*

Proof. Let $M \leq A_5$ be maximal. Note that a subgroup H of index n in A_5 induces a non-trivial homomorphism $A_5 \rightarrow S_n$ (via the action of A_5 on the cosets of H), and by simplicity of A_5 this map must be an injection. Hence we must have $n \geq 5$, i.e. $o(H) \leq 12$.

Now, any subgroup with a fixed point is contained in a copy of A_4 , and each copy of A_4 is maximal in A_5 (since $o(A_4) = 12$). So if M has a fixed point, $M \cong A_4$.

Next, say M acts transitively. Then we must have $5|o(M)$, so $o(M)$ is either five or ten. Now, $(15)(24)$ normalizes $\langle(12345)\rangle$, so $N(\langle(12345)\rangle) \neq A_5$ has order at least ten, and in fact its order can be no larger. Thus $N(\langle(12345)\rangle) \cong D_{10}$, and hence every subgroup of order five is contained in a copy of D_{10} (subgroups of order five are Sylow subgroups, hence conjugate). So if M acts transitively, $M \cong D_{10}$.

Finally, if M neither acts transitively nor has a fixed point, then the orbits of $\{1, 2, 3, 4, 5\}$ under the action of M must have sizes two and three. In this

case, one may check that the elements of A_5 permuting the sets $\{1, 2\}$ and $\{3, 4, 5\}$ amongst themselves form a subgroup isomorphic to S_3 (in fact, this is the standard embedding of S_{n-2} in A_n). This subgroup is in fact maximal in A_5 , since any larger subgroup would be transitive and therefore have order divisible by 5. So in this last case, $M \cong S_3$.

Now we turn to the number of subgroups of each type. Any copy of A_4 is maximal since it has order twelve, and thus must be one of the obvious five copies. Any copy of D_{10} certainly normalizes a $\mathbb{Z}/5$, and so the only such subgroups are the normalizers of the Sylow 5-subgroups of A_5 . There are six such subgroups, as is easily checked. Finally, we claim that any copy of S_3 is maximal (and hence arises in the manner described above). If $H \leq A_5$ were isomorphic to S_3 but not maximal, then H would be contained in a copy of A_4 . Thus it sufficed to check that A_4 has no subgroup of order six. Any such subgroup would be normal, and hence would have to contain all the Sylow 3-subgroups of A_4 . But there are four of these, totalling nine elements. So all the S_3 arise in the above manner, and are clearly conjugate in S_5 . $\square$

Corollary 4.2.3 *All chains in $\mathcal{C} = \mathcal{C}(A_5)$ of length greater than three are of the form $\{x\} \leq xC_2 \leq xD_4 \leq xA_4$ (where C_2 denotes the cyclic group of order two).*

Proof. Underlying every chain in $\mathcal{C}$ is a chain of the same length in $S(A_5) \cup \{1\}$. These latter chains each end in either a copy of S_3 , a copy of D_{10} , or a copy of A_4 . Since the orders of S_3 and D_{10} are products of two distinct primes, the longest chains ending at these subgroups are of length three. So we only need to consider chains ending at A_4 . The subgroups of A_4 are of orders two, three and four (there are no subgroups of order six, as discussed earlier). So any chain of length four must be of the form described in the corollary (noting that the unique Sylow 2-subgroup of A_4 is isomorphic to D_4). $\square$

Claim 4.2.4 *Let $\bar{\mathcal{C}}$ denote the poset $\mathcal{C}$ with all cosets of all copies of D_4 removed. Then $\Delta(\bar{\mathcal{C}})$ is two-dimensional and $\bar{\mathcal{C}} \simeq \mathcal{C}$.*

Proof. The fact that $\Delta(\bar{\mathcal{C}})$ is two-dimensional follows immediately from Corollary 4.2.3. To show that $\bar{\mathcal{C}} \simeq \mathcal{C}$, we apply Quillen's Theorem to the inclusion $i : \bar{\mathcal{C}} \rightarrow \mathcal{C}$. If $H < A_5$ is not isomorphic to D_4 , then $i^{-1}(\mathcal{C}_{\geq xH})$ is a cone on xH for any $x \in A_5$. When $H \cong D_4$, we have $i^{-1}(\mathcal{C}_{\geq xH}) = \{xK :$

$H < K < G\}$. Any such K is isomorphic to A_4 , and each D_4 lies in a unique A_4 (its normalizer in A_5). So $i^{-1}(\mathcal{C}_{\geq xH})$ is a single point, and thus Quillen's Theorem shows that i is a homotopy equivalence. $\square$

We now turn to simple connectivity. First we will show that A_5 satisfies the conditions of Theorem 3.1.9, thereby showing $\mathcal{C}(A_5)$ is simply connected. Then we will give a second, more conceptual proof using 2-transitive actions of A_5 .

Claim 4.2.5 *For each pair $x, y \in A_5$ with $\langle x, y \rangle = A_5$, there is an element $z \in A_5$ such that $\langle x, z \rangle$, $\langle y, z \rangle$, and $\langle zx, zy \rangle$ are all proper subgroups. Theorem 3.1.9 now shows that $\mathcal{C}(A_5)$ is simply connected.*

Proof. We will prove this result by considering representatives of all possible automorphism classes of generating pairs (i.e. all sets of the form $\{(\phi(x), \phi(y)) : \phi \in \text{Aut}(A_5), \langle x, y \rangle = A_5\}$) and finding, for each such representative pair, an appropriate element z (in some cases it is not actually necessary to choose a specific representative). Luckily the total number of automorphism classes is relatively small. We will break the problem down by considering the orders of x and y .

- **$\mathbf{o}(x) = \mathbf{o}(y) = 2$:** In any group, the subgroup generated by two elements of order two is dihedral, and in particular two such elements cannot generate a simple group. (The latter statement is easily checked by verifying that $x, y \in N(\langle xy \rangle)$, which is a proper subgroup by simplicity.)
- **$\mathbf{o}(x) = 2, \mathbf{o}(y) = 3$:** Since x and y generate A_5 they cannot have a common fixed point. Up to automorphism, we may assume that $x = (12)(45)$. The three-cycle y must move an element from each two-cycle, since otherwise x and y would sit in a copy of S_3 . So up to automorphism, we may assume that $y = (234)$. Now, setting $z = (123)$ we see that $\langle x, z \rangle$ lies a copy of S_3 (its action has orbits $\{1, 2, 3\}, \{4, 5\}$) and $\langle y, z \rangle$ fixes 5 and is contained in an A_4 . Furthermore, zx and zy have order 2, so $\langle zx, zy \rangle \neq A_5$ by the above discussion.
- **$\mathbf{o}(x) = 2, \mathbf{o}(y) = 5$:** First, x lies in a copy of A_4 and y lies in a copy of D_{10} . These subgroups must intersect nontrivially in A_5 , since otherwise we would have $o(A_5) \geq o(A_4)o(D_{10}) = 120$, a contradiction.

Since the only common divisor of $o(D_{10})$ and $o(A_4)$ is two, there must be an element z of order two in this intersection. (Note that x does not lie in the intersection, since $\langle x, y \rangle = A_5$.) By choice of z , we see that $\langle z, x \rangle$ and $\langle z, y \rangle$ are proper subgroups. In addition, since D_{10} is dihedral the product of an element of order two and an element of order five has order two, i.e. $o(zy) = 2$. Similarly the product of two elements of order two in A_4 has order two, so $o(zx) = 2$. So once again we see that $\langle zx, zy \rangle \neq A_5$.

- **$o(\mathbf{x}) = o(\mathbf{y}) = 3$** Let $x = (123)$. In order for x and y to generate, y must move both 4 and 5, and up to automorphism we may assume that $y = (345)$. Let $z = (234)$. Then x and z fix 5, and y and z fix 1, so neither pair generates A_5 . Once again we find that $o(zx) = o(zy) = 2$.
- **$o(\mathbf{x}) = 3, o(\mathbf{y}) = 5$** : Let $x = (123)$. It is easily checked that there is an element $z = (ab)(45) \in N(\langle y \rangle)$, with $a, b \in \{1, 2, 3\}$. Now, x and z each lie in the copy of S_3 with orbits $\{1, 2, 3\}$ and $\{4, 5\}$, and hence do not generate A_5 . As discussed above, $o(zy) = 2$ and since zx contains the transposition (45) it has order two as well.
- **$o(\mathbf{x}) = o(\mathbf{y}) = 5$** : The normalizers of $\langle x \rangle$ and $\langle y \rangle$ must intersect non-trivially, since otherwise we would have $o(A_5) \geq o(D_{10})^2$, a contradiction. Since x and y generate, $\langle x \rangle \neq \langle y \rangle$ and hence $\langle x \rangle \cap \langle y \rangle = \{1\}$. This implies that the normalizers of $\langle x \rangle$ and $\langle y \rangle$ meet in a subgroup of order two. If z is the non-trivial element in this intersection, then we see that $\langle x, z \rangle$ and $\langle y, z \rangle$ are proper and $o(zx) = o(zy) = 2$.

This covers all possible generating pairs, and thus completes the proof. $\square$

Alternate Proof of Simple Connectivity. In the above proof, it was shown that for each edge $\{x, y\} \in \mathcal{M}^2(A_5)$ with $o(x) = 2$, there is an element $z \in G$ satisfying the conditions of Theorem 3.1.9. Consider the presentation for $\pi_1(\mathcal{M}(A_5))$ given by Theorem 2.1.9, using as a maximal tree all edges $\{1, g\}$, $g \in G$ ($g \neq 1$). Note that this is the same presentation as was used in the proof of Theorem 3.1.9, and recalling that proof, the existence of the element z implies that the generators $(x, y) = (y, x)^{-1}$ are trivial in $\pi_1(\mathcal{M}(A_5))$.

Starting from this fact (that generators (x, y) with $o(x) = 2$ are trivial) we will complete the proof in a less ad hoc manner. The same method will be used in the next section to show that $\mathcal{C}(PSL_2(\mathbb{F}_7))$ is simply connected.

Let $\{g, h\}$ be any edge of $\mathcal{M}^2(A_5)$. We must show that the corresponding generators of $\pi_1(\mathcal{M}(A_5))$ are trivial. It will suffice to show that there exists an element z of order two and a subgroup $K \in \mathcal{C}(A_5)$ s.t. $h \equiv g \equiv z \pmod{K}$. This is true because then the set $\{g, h, z\}$ forms a 2-simplex in $\mathcal{M}(A_5)$ and then the relations in Theorem 2.1.9 show that $(g, h)(h, z)(z, g) = 1$. Since $o(z) = 2$, we have $(h, z) = (z, g) = 1$, so $(g, h) = 1$ as desired.

Now, say G acts 2-transitively on the set X , and assume that some element z with $o(z) = 2$ acts non-trivially. Let $H = \text{Stab}(x)$ for some $x \in X$, and note that for any $y \in X$, the set $S_{x,y} = \{g \in G : g \cdot x = y\}$ is just the coset kH , where $k \cdot x = y$. Now, since z acts non-trivially and the action is 2-transitive, some conjugate of z sends x to y , i.e. there is an element of order two in every non-trivial coset of H . Thus we can prove simple connectivity by showing that every two-element set $\{g, h\} \subset G$ lies in a coset kH where G acts 2-transitively on some set X , H acts as the stabilizer of a point, and some element of order two acts non-trivially.

First, consider the standard action of A_5 on the set $\{1, 2, 3, 4, 5\}$. This action is clearly 2-transitive, and the stabilizer of a point is isomorphic to A_4 . Every non-trivial element acts non-trivially, so certainly elements of order two act non-trivially.

Next, consider the action of A_5 on its Sylow 5-subgroups (by conjugation). Call this set S . There are six subgroups in S , and each of them acts transitively on the other five (certainly each acts non-trivially, and since the orbits must have order dividing 5, the action is transitive). Now given any $H_1 \neq K_1$, $H_2 \neq K_2 \in S$, there is an element $y \in H_1$ with $K_1^y = K_2$ and there is an element $z \in K_2$ with $H_1^z = H_2$. We have $H_1^{yz} = H_1^z = H_2$ and $K_1^{yz} = K_2^z = K_2$, showing that the action is indeed 2-transitive. Note that the stabilizer of a subgroup in S is just its normalizer, a copy of D_{10} .

We claim that every two-element set $\{g, h\}$ lies in either a coset of a copy of A_4 or in a coset of a copy of D_{10} . In other words, we claim that $g^{-1}h$ always lies in a subgroup isomorphic to either A_4 or D_{10} . But every cyclic subgroup of A_5 has order two, three, or five, so this follows immediately. $\square$

4.3 The Coset Poset of $PSL_2(\mathbb{F}_7)$

In this section we will consider the finite simple group $G = PSL_2(\mathbb{F}_7)$. This group has order 168, and is the next larger simple group after A_5 . Our goal is to show that $\mathcal{C}(G)$ is simply connected. Unfortunately, we do not know

whether $\mathcal{C}(G)$ is (homotopy equivalent to) a bouquet of spheres or something more exotic.

The proof of simply connectivity will require a careful understanding of the subgroup poset of G , and in particular we will need to compute the Möbius function of G (defined in Section 4.3.2). Rather than restrict ourselves to the case $p = 7$, we will give a detailed description of the subgroups of $PSL_2(\mathbb{F}_p)$ for all primes $p \geq 5$. This computation will be based on Burnside's exposition in [8]. The computation of the Möbius function of G will be based on the (brief) calculation given by Hall in his original paper [11] on Möbius inversion.

The proof of simple connectivity will be analogous to the second proof for A_5 (given in the previous section), and will utilize 2-transitive actions of G . The Möbius function of G will be used to count automorphism classes of generating pairs. In the case of A_5 this was unnecessary simply because these classes were easy to enumerate by hand.

Once we have proven simple connectivity, we will briefly discuss other facts about the homotopy-type of $\mathcal{C}(G)$.

We will begin by defining the groups we propose to study, and then describing a useful permutation representation for them.

Fix a prime p , and let $\mathbb{F}_p \cong \mathbb{Z}/p$ denote the finite field with p elements. We define $GL_n(\mathbb{F}_p)$, the n -dimensional general linear group over $\mathbb{F}_p$, to be the multiplicative group of invertible $n \times n$ matrices with entries in $\mathbb{F}_p$, and we define the n -dimensional special linear group, $SL_n(\mathbb{F}_p) \leq GL_n(\mathbb{F}_p)$, to be the subgroup of matrices with determinant one.

Let $\Gamma_{n,p}$ denote the center of $SL_n(\mathbb{F}_p)$. The n -dimensional projective special linear group, $PSL_n(\mathbb{F}_p)$, is defined to be the quotient $SL_n(\mathbb{F}_p)/\Gamma_{n,p}$. The following theorem is well-known (see, for example, [19, Theorem 3.2.9]).

Theorem 4.3.1 *If $n > 2$ or $p > 3$, then $PSL_n(\mathbb{F}_p)$ is a simple group.*

We will mainly be interested in the cases where $n = 2$ and $p > 3$. In these cases, direct computation shows that $\Gamma_{2,p} = \{I, -I\}$, so $o(PSL_2(\mathbb{F}_p)) = \frac{1}{2}o(SL_2(\mathbb{F}_p))$. Our first goal is to determine the order of $PSL_2(\mathbb{F}_p)$.

Claim 4.3.2 *If $p > 3$, then $o(PSL_2(\mathbb{F}_p)) = \frac{1}{2}p(p^2 - 1)$*

Proof. An element of $GL_2(\mathbb{F}_p)$ may be thought of as a pair (v, w) where v and w are linearly independent vectors in $\mathbb{F}_p^2$. Thus in forming an element

of $GL_2(\mathbb{F}_p)$, we have $p^2 - 1$ choices for v (v must simply be non-zero), and then $p^2 - p$ choices for w (w must not lie in the span of v). Thus we have $o(GL_2(\mathbb{F}_p)) = (p^2 - 1)(p^2 - p)$.

The exact sequence

$$1 \longrightarrow SL_2(\mathbb{F}_p) \hookrightarrow GL_2(\mathbb{F}_p) \xrightarrow{\det} \mathbb{F}_p^* \longrightarrow 1$$

(where $\det$ denotes the determinant homomorphism) shows that

$$o(SL_2(\mathbb{F}_p)) = \frac{(p^2 - 1)(p^2 - p)}{p - 1} = (p + 1)(p^2 - p) = p(p^2 - 1).$$

As mentioned above, $PSL_2(\mathbb{F}_p)$ has index two in $SL_2(\mathbb{F}_p)$, completing the proof. $\square$

From now on we will write $o(PSL_2(\mathbb{F}_p)) = 2pqr$, where $q = \frac{p-1}{2}$ and $r = \frac{p+1}{2}$. The main advantage of this notation is the fact that p , q and r are pair-wise relatively prime.

We will now describe a useful action of $PSL_2(\mathbb{F}_p)$. Let $V = \mathbb{F}_p \oplus \mathbb{F}_p$ be a two-dimensional vector space over $\mathbb{F}_p$ and define an equivalence relation $\sim$ on $V - \{0\}$ by setting $v \sim \lambda v$ for all $v \in V - \{0\}$, $\lambda \in \mathbb{F}_p^*$. We identify the set of equivalence classes with the projective line $\mathbb{F}_p \cup \{\infty\}$ (where ∞ is just a formal symbol) via the map $(v_1, v_2) \mapsto \frac{v_1}{v_2}$, with the convention that $\frac{a}{0} = \infty$ for any $a \in \mathbb{F}_p$ (this map is clearly well-defined on equivalence classes, and it is not hard to check that it is a bijection).

Then $GL_2(\mathbb{F}_p)$ acts on $V - \{0\} / \sim$ (and hence on $\mathbb{F}_p \cup \{\infty\}$) by multiplication. Restricting to $SL_2(\mathbb{F}_p)$, it is clear that this action factors through the quotient map $\pi : SL_2(\mathbb{F}_p) \rightarrow PSL_2(\mathbb{F}_p)$, and we obtain an action of $PSL_2(\mathbb{F}_p)$ on $\mathbb{F}_p \cup \{\infty\}$.

It will be convenient to describe another way of writing elements in $PSL_2(\mathbb{F}_p)$. Consider the group M_p of (invertible) Möbius transformations $\mathbb{F}_p \cup \{\infty\} \rightarrow \mathbb{F}_p \cup \{\infty\}$. These are maps of the form $x \mapsto \frac{ax+b}{cx+d}$ (where $x \in \mathbb{F}_p \cup \{\infty\}$ and $a, b, c, d \in \mathbb{F}_p$) with $ad - bc = 1$, and multiplication is defined by composition. We use the conventions $\frac{a\infty+b}{c\infty+d} = \frac{a}{c}$ and $\frac{z}{0} = \infty$ (in all other situations, we simply add and multiply in $\mathbb{F}_p$). We write elements of M_p in the form $\frac{ax+b}{cx+d}$, where x is an indeterminate, and it is easy to check that the inverse of $\frac{ax+b}{cx+d} \in M_p$ is $\frac{dx-b}{-cx+a}$. Since these maps are invertible, they

are bijections and hence M_p acts on $\mathbb{F}_p \cup \{\infty\}$. Note that in this notation the identity element of M_p is simply x , but to avoid confusion we will sometimes denote the identity of M_p by 1 or by id_M .

It is easy to check that the obvious map $SL_2(\mathbb{F}_p) \longrightarrow M_p$ is a homomorphism with kernel $\{I, -I\}$, and thus induces a (canonical) isomorphism $PSL_2(\mathbb{F}_p) \longrightarrow M_p$. In addition, this isomorphism commutes with the actions on $\mathbb{F}_p \cup \{\infty\}$. It is easy to see that the action of M_p on $\mathbb{F}_p \cup \{\infty\}$ is transitive, and we will later see that it is in fact 2-transitive.

4.3.1 The Subgroups of $PSL_2(\mathbb{F}_p)$

We will now describe the subgroups of $PSL_2(\mathbb{F}_p) \cong M_p$, $p > 5$. We will refer mainly to $M = M_p$ during this analysis. Our first goal will be to prove the following theorem describing the cyclic subgroups of M and their normalizers. The proof will be broken down into a number of claims.

Theorem 4.3.3 *The set of maximal cyclic subgroups of M consists of:*

1. $p+1$ cyclic subgroups of order p , each fixing a single point of $\mathbb{F}_p \cup \{\infty\}$;
2. pr cyclic subgroups of order q , each fixing two points of $\mathbb{F}_p \cup \{\infty\}$;
3. pq cyclic subgroups of order r , each acting freely on $\mathbb{F}_p \cup \{\infty\}$.

Each of the three collections above forms a conjugacy class in M . In addition, the intersection of any two of the subgroups listed is trivial, so each element of M lies in a unique maximal cyclic subgroup.

Any cyclic subgroup of order p has normalizer isomorphic to $\mathbb{Z}/p \rtimes \mathbb{Z}/q$ (for some action of $\mathbb{Z}/q$ on $\mathbb{Z}/p$, which we will not need to specify).

The normalizer of any other (non-trivial) cyclic subgroup $C \leq M$ is isomorphic to either D_{2q} or D_{2r} , depending on whether $o(C)$ divides q or r (we use the convention $D_{2n} = \mathbb{Z}/n \rtimes \mathbb{Z}/2$, where the generator of $\mathbb{Z}/2$ acts by inversion).

Note that by “maximal cyclic subgroup” we mean a cyclic subgroup not contained in any larger cyclic subgroup. Also, recall that p , q and r are pairwise relatively prime, so in showing that none of the cyclic subgroups listed above intersect non-trivially, we need only consider intersections between groups of the same order.

Before beginning the proof, we record two useful lemmas.

Lemma 4.3.4 *Let G be a finite group and $H \leq G$ a subgroup. Then the number of conjugates of H in G is equal to $(G : N_G(H))$, the index of the normalizer of H .*

This result follows easily by considering the actions of G on the cosets of $N(H)$ (by left multiplication) and on the conjugates of H (by conjugation).

Lemma 4.3.5 *Let G be a finite group and let $H \leq G$ be a cyclic subgroup. Assume that any two conjugates of H intersect trivially. Then for any $K \leq H$, we have $N_G(K) = N_G(H)$.*

Proof. Since K is the unique subgroup of its order in H , we have $K \triangleleft N_G(H)$. In addition, we see that K has the same number of conjugates as H , since any two conjugates of H intersect trivially. Letting c_H and c_K denote the number of conjugates of H and K in G , Lemma 4.3.4 gives

$$o(N_G(K)) = \frac{o(G)}{c_K} = \frac{o(G)}{c_H} = o(N_G(H)),$$

completing the proof. $\square$

We will now examine the subgroups of order p in M . Note that these are in fact Sylow p -subgroups, and hence form a single conjugacy class in M . Any two of intersect trivially, since they are of prime order.

Claim 4.3.6 *There are $p + 1$ cyclic subgroups of order p in M , each with normalizer isomorphic to $\mathbb{Z}/p \rtimes \mathbb{Z}/q$.*

Proof. Consider the element $x + 1 \in M$. Note that $(x + 1)^n = x + n$, and hence $o(x + 1) = p$. Thus $\langle x + 1 \rangle$ is cyclic subgroup of order p , and note that this subgroup fixes $\infty \in \mathbb{F}_p \cup \{\infty\}$ but does not fix any other point.

If $m \in M$, then $m\langle x + 1 \rangle m^{-1}$ will fix the point $m(\infty)$. Thus $N(\langle x + 1 \rangle) \leq \text{Stab}(\infty)$, since $m \notin \text{Stab}(\infty)$ implies $m\langle x + 1 \rangle m^{-1}$ does not fix ∞ . There are $p + 1$ points in $\mathbb{F}_p \cup \{\infty\}$, giving $p + 1$ distinct conjugates of $\langle x + 1 \rangle$ (recall that the action of M on $\mathbb{F}_p \cup \{\infty\}$ is transitive).

To complete the proof it will suffice to show that $\text{Stab}(\infty) \simeq \mathbb{Z}/p \rtimes \mathbb{Z}/q$. Since $\frac{a\infty+b}{c\infty+d} = \frac{a}{c}$, we see that

$$\text{Stab}(\infty) = \left\{ \frac{ax+b}{cx+d} \in M : c = 0 \right\} = \left\{ \frac{ax+b}{a^{-1}} \right\}.$$

Since we must have $a \neq 0$, there are $\frac{1}{2}(p-1)p = pq$ elements in this subgroup. Note that $\langle x+1 \rangle$ is a Sylow p -subgroup of $\text{Stab}(\infty)$, and thus is either normal or has at least $p+1$ conjugates within $\text{Stab}(\infty)$. These $p+1$ conjugates would contain a total of $(p+1)(p-1) + 1 > pq = o(\text{Stab}(\infty))$ elements, and thus we have $\langle x+1 \rangle \triangleleft \text{Stab}(\infty)$.

Finally, note that there is a copy of $\mathbb{Z}/q$ in $\text{Stab}(\infty)$, generated by the element $\frac{zx}{z-1}$ with z a generator of $\mathbb{F}_p^*$. (Since $\left(\frac{zx}{z-1}\right)^n = \frac{z^n x}{z^n - 1}$, we have $\frac{z^n x}{z^n - 1} = x = id_M$ iff $z^n = z^{-n}$, i.e. iff $z^{2n} = 1$. Now, $o(z) = o(\mathbb{F}_p^*) = p-1$, so $o\left(\frac{zx}{z-1}\right) = \frac{p-1}{2} = q$.) Thus we have found a complement (in $\text{Stab}(\infty)$) to the normal subgroup $\langle x+1 \rangle$, and $\text{Stab}(\infty) \cong \mathbb{Z}/p \rtimes \mathbb{Z}/q$. $\square$

We now turn to the cyclic subgroups of order q . We have already exhibited one such subgroup $\langle \frac{zx}{z-1} \rangle$ for z a generator of $\mathbb{F}_p^*$ in the proof of Claim 4.3.6, and the other such subgroups are conjugates of this one. First we will need two simple lemmas about the action of M on $\mathbb{F}_p \cup \{\infty\}$.

Lemma 4.3.7 *The group M acts 2-transitively on $\mathbb{F}_p \cup \{\infty\}$.*

Proof. Note that, as shown in the proof of Claim 4.3.6, for each $x \in \mathbb{F}_p \cup \{\infty\}$, there is a subgroup P_x of order p in $\text{Stab}(x)$. Since no element of M other than the identity acts trivially on $\mathbb{F}_p \cup \{\infty\}$, we see that P_x acts transitively on $\mathbb{F}_p \cup \{\infty\} - \{x\}$ (because this is a set of cardinality p). Now, given any $a \neq b, c \neq d \in \mathbb{F}_p \cup \{\infty\}$, there is an element $y \in P_a$ with $y(b) = d$ and there is an element $z \in P_d$ with $z(a) = c$. We have $zy(a) = z(a) = c$ and $zy(b) = z(d) = d$, showing that M is indeed 2-transitive. $\square$

Lemma 4.3.8 *No non-trivial element of M can fix more than two points in $\mathbb{F}_p \cup \{\infty\}$.*

Proof. Notice that fixed points (other than ∞) correspond to solutions of a quadratic polynomial: if $m = \frac{ax+b}{cx+d} \in M$, then

$$m(x) = x \iff ax + b = cx^2 + dx \iff cx^2 + (d-a)x - b = 0.$$

If m has more than two fixed points in $\mathbb{F}_p$, this polynomial has more than two roots and must be zero. Thus m fixes every point of $\mathbb{F}_p \cup \{\infty\}$, i.e. m is the identity.

Finally, if $m(\infty) = \infty$, then $m = \frac{ax+b}{d}$ and any other fixed point x satisfies $\frac{ax+b}{d} = x$, i.e. $(a-d)x - b = 0$, and if $m \neq 1$ this linear equation can have at most one root. $\square$

For $x, y \in \mathbb{F}_p \cup \{\infty\}$, let $\text{Stab}(x, y) = \text{Stab}(x) \cap \text{Stab}(y)$.

Claim 4.3.9 *For any two distinct points $x, y \in \mathbb{F}_p \cup \{\infty\}$, $\text{Stab}(x, y) \cong \mathbb{Z}/q$. This yields pr conjugate copies of $\mathbb{Z}/q$, and any two intersect trivially. Furthermore, if $t \in \text{Stab}(x, y)$ ($x \neq y$), then $N(\langle t \rangle) = N(\text{Stab}(x, y)) \cong D_{2q}$.*

Proof. It is easily checked that $\text{Stab}(0, \infty) = \left\{ \frac{ax}{a-1} : a \in \mathbb{F}_p \right\}$, and this subgroup is generated by any element $\frac{zx}{z-1}$ with z a generator of $\mathbb{F}_p$. As was shown in the proof of Claim 4.3.6, such an element has order q . Since M is 2-transitive, $\text{Stab}(x, y) \cong \text{Stab}(0, \infty)$ for any $x \neq y \in \mathbb{F}_p \cup \{\infty\}$. There are $\binom{p+1}{2} = pr$ such stabilizers, and two of them intersect trivially by Lemma 4.3.8.

Finally, we must consider the normalizers of these subgroups. Clearly, $\text{Stab}(0, \infty) \leq \text{Stab}(\{0, \infty\})$, where the latter denotes the elements of M which fix $\{0, \infty\}$ as a set. We claim that $\text{Stab}(\{0, \infty\}) = N(\text{Stab}(0, \infty))$ and that $\text{Stab}(\{0, \infty\}) \cong D_{2q}$.

Any element $m \in \text{Stab}(\{0, \infty\})$ either fixes both points or swaps them. In the former case $m \in \text{Stab}(0, \infty)$ and in the latter we have $m = \frac{b}{-b^{-1}x}$ for some $b \in \mathbb{F}_p^*$. In either case we obtain q elements, so $o(\text{Stab}(\{0, \infty\})) = 2q$ and hence

$$\text{Stab}(0, \infty) \triangleleft \text{Stab}(\{0, \infty\}).$$

In addition, for any $b \in \mathbb{F}_p^*$ we have

$$\left(\frac{b}{-b^{-1}x} \right)^2 = \frac{b}{-b^{-1} \frac{b}{-b^{-1}x}} = x = id_M,$$

so all elements in $\text{Stab}(\{0, \infty\}) - \text{Stab}(0, \infty)$ have order two. To show that $\text{Stab}(\{0, \infty\})$ is dihedral, all that remains to be checked is that an element in $\text{Stab}(\{0, \infty\}) - \text{Stab}(0, \infty)$ conjugates any element of $\text{Stab}(0, \infty)$ to its inverse. This is easily verified by multiplying representative matrices in $SL_2(\mathbb{F}_p)$. Lemmas 4.3.4 and 4.3.5 complete the proof. $\square$

We now come to the cyclic subgroups of order r . These arise in a rather different manner, since they act freely on $\mathbb{F}_p \cup \{\infty\}$.

Let $K = \mathbb{F}_{p^2}$, and fix a basis β for K over $k = \mathbb{F}_p$. For $z \in K^*$, let $L_z \in GL_2(\mathbb{F}_p)$ denote the matrix (in the basis β) corresponding to the map $y \mapsto zy$ ($y \in K$), and let $H = \{L_z : z \in K^*\} \leq GL_2(\mathbb{F}_p)$. Let $\tilde{H} = \phi(H \cap SL_2(\mathbb{F}_p))$, where $\phi : SL_2(\mathbb{F}_p) \rightarrow M$ denotes the obvious map. We will show that $\tilde{H} \cong \mathbb{Z}/r$.

We now record two useful lemmas about the Galois extension K/k .

Lemma 4.3.10 *Every element of k has a square root in K .*

Proof. This follows from the fact that any degree two extension of k is isomorphic to K , but can also be seen directly, as follows. The (multiplicative) group K^* is isomorphic to the (additive) group $\mathbb{Z}/(p^2 - 1)$, and squares in K^* correspond to even residues in $\mathbb{Z}/(p^2 - 1)$. The subgroup $k^* \leq K^*$ has order $p - 1$, and maps to $\langle p + 1 \rangle \leq \mathbb{Z}/(p^2 - 1)$. Since $p + 1$ is even, these elements all have “square roots” in $\mathbb{Z}/(p^2 - 1)$ and thus all elements of k^* have square roots in K^* . $\square$

Lemma 4.3.11 *Let $\sigma : K \rightarrow K$ denote the p^{th} -power map, i.e. $\sigma(z) = z^p$. Then $\text{Gal}(K/k) = \{I, \sigma\}$, and $\det(\sigma) = -1$.*

Proof. First, the fact that σ is a field automorphism of K follows from the binomial formula

$$(x + y)^p = \sum_{i=0}^p \binom{p}{i} x^i y^{p-i},$$

since for $i \neq 0, p$ we have $p \mid \binom{p}{i} = \frac{p!}{i!(p-i)!}$. Next, Fermat’s Little Theorem shows that for any $z \in k$, $z^p = z$ and hence σ is an automorphism of K/k . Finally, since $\deg(K/k) = 2$, $\text{Gal}(K/k) \cong \mathbb{Z}/2$ and hence $\text{Gal}(K/k) = \{I, \sigma\}$.

Finally we must compute $\det(\sigma)$. (Note that an element of the Galois group fixes k and thus is also in $GL_2(\mathbb{F}_p)$, so $\det(\sigma)$ makes sense). By the Normal Basis Theorem, there is an element $y \in K$ such that $\{y, \sigma(y)\}$ is a basis for K over k . Since σ has order two, it interchanges these basis vectors and thus has determinant -1 . $\square$

Claim 4.3.12 *The subgroup $\tilde{H}$ has pq conjugates in M . Any two of these subgroups intersect trivially, and we have:*

1. $\tilde{H} \cong \mathbb{Z}/r$;
2. $\tilde{H}$ acts freely on $\mathbb{F}_p \cup \{\infty\}$;
3. For any non-trivial subgroup $T \leq \tilde{H}$, $N(T) \cong D_{2r}$.

Proof. First note that $\tilde{H}$ is cyclic because K^* is cyclic. To determine the order of $\tilde{H}$ we must first determine $o(H \cap SL_2(\mathbb{F}_p))$, i.e. we must determine which elements of H have determinant one.

It is a result in Galois theory that for any $z \in K^*$, $\det(L_z) = N_{K/k}(z)$ where $N_{K/k}(z)$ denotes the norm of z , i.e. the product of its Galois conjugates in K . Lemma 4.3.11 now shows that $\det(L_z) = z\sigma(z) = z^{p+1}$. Thus $H \cap SL_2(\mathbb{F}_p) \cong \{z \in K^* : z^{p+1} = 1\} \cong \{a \in \mathbb{Z}/(p^2 - 1) : (p+1)a = 0\}$, and the last set clearly has cardinality $p+1$. Finally, since $L_{-1} = -I$,

$$o(\tilde{H}) = \frac{1}{2}o(H \cap SL_2(\mathbb{F}_p)) = \frac{p+1}{2} = r.$$

Next we consider the action of $\tilde{H}$ on $\mathbb{F}_p \cup \{\infty\}$. We want to show that this action is free, i.e. that no element of $\tilde{H}$ fixes a point on $\mathbb{F}_p \cup \{\infty\}$. Consider the action of $H \cap SL_2(\mathbb{F}_p)$ on $\mathbb{F}_p \cup \{\infty\}$. If $L_y \in H \cap SL_2(\mathbb{F}_p)$ fixes a point in $\mathbb{F}_p \cup \{\infty\}$, then $L_y(v) = yv = \lambda v$ for some $v \in K - \{0\}$ and some $\lambda \in k$. But then $y = \lambda$, i.e. $y \in k$, and we have

$$1 = \det(L_y) = N(y) = y\sigma(y) = y^2$$

(since σ fixes k). Hence $y = \pm 1$ and $\phi(L_y) = \phi(-I) = id_M$. So $\tilde{H}$ acts freely on $\mathbb{F}_p \cup \{\infty\}$.

We now consider $N_M(\tilde{H})$. Let $y \in K^*$ be an element with $\det(L_y) = -1$. (Such elements exist, since under the isomorphism $K^* \xrightarrow{\cong} \mathbb{Z}/(p^2 - 1)$ they correspond to elements $t \in \mathbb{Z}/(p^2 - 1)$ with $(p+1)t = \frac{p^2-1}{2}$.) By Lemma 4.3.11 we have $\det(\sigma L_y) = (-1)^2 = 1$, so $\sigma L_y \in SL_2(\mathbb{F}_p)$. Let $\alpha = \phi(\sigma L_y)$. We claim that $\langle \tilde{H}, \alpha \rangle \cong D_{2r}$.

To prove this, it suffices to show that $o(\alpha) = 2$ and that for any $h \in \tilde{H}$, $\alpha h \alpha^{-1} = h^{-1}$. First, if $v \in K$ then

$$(\sigma L_y)^2 v = (\sigma L_y \sigma L_y) v = \sigma(y\sigma(yv)) = \sigma(y)yv = \det(y)v = -v$$

and hence $(\sigma L_y)^2 = -I$ and $\alpha^2 = id_M$. Thus $o(\alpha) \leq 2$, and we will see that α is non-trivial because its action on $\tilde{H}$ is non-trivial.

Next we wish to prove that $\alpha h \alpha^{-1} = h^{-1}$ for any $h \in \tilde{H}$. Choose $z \in K^*$ with $\phi(L_z) = h$. Then for any $v \in K$ we have

$$\begin{aligned} (\sigma L_y) L_z (\sigma L_y)^{-1} v &= (\sigma L_y L_z L_y^{-1} \sigma) v = \sigma(yzy^{-1}\sigma(v)) \\ &= \sigma(z\sigma(v)) = \sigma(z)v = z^{-1}v = L_{z^{-1}}v, \end{aligned}$$

where the next to last equality follows because $1 = \det(L_z) = z\sigma(z) \implies \sigma(z) = z^{-1}$. So $(\sigma L_y) L_z (\sigma L_y)^{-1} = L_{z^{-1}}$ and hence $\alpha h \alpha^{-1} = h^{-1}$, as desired.

By Lemma 4.3.4, the proof will be complete once we show that $\tilde{H}$ has pq conjugates, any two of which intersect trivially. We will describe pq different bases for K over k , and changing basis in $GL_2(\mathbb{F}_p)$ will yield pq distinct conjugates of $\tilde{H}$. Consider all bases of the form $\beta_{y,l} = \{\sqrt{y} + l, 1\}$ where $y, l \in \mathbb{F}_p$ and y is not a square in $\mathbb{F}_p$ (there is ambiguity in the symbol $\sqrt{y}$ since there are two roots in K of the equation $x^2 = y$; for each y we arbitrarily choose one of these roots and denote it by $\sqrt{y}$). Each of these is clearly a basis (since $\sqrt{y} \notin \mathbb{F}_p$), and there are pq of them since we have $\frac{p-1}{2} = q$ choices for y and p choices for l .

Let $H^{\beta_{y,l}}$ denote the conjugate of H obtained by changing basis from β to $\beta_{y,l}$. By writing out the matrices involved, one finds that

$$H^{\beta_{y,l}} \cap H^{\beta_{z,n}} \cap SL_2(\mathbb{F}_p) = \{I, -I\}$$

unless $y = z$ and $l = n$. This implies that the resulting conjugates of $\tilde{H}$ have trivial intersection in M . $\square$

To complete the proof of Theorem 4.3.3, we must simply show that each element of M lies in one of the cyclic subgroups we have described. This follows from a simple counting argument: the total number of elements involved in these subgroups is

$$\begin{aligned} (p+1)(p-1) + pr(q-1) + pq(r-1) + 1 &= p^2 + 2pqr - pr - pq \\ &= 2pqr + p^2 - \frac{p(p+1)}{2} - \frac{p(p-1)}{2} = 2pqr = o(M). \end{aligned}$$

Our next goal will be to study subgroups $Q \leq M$ with $Q \cong D_4$. Following Burnside, we call such subgroups *quadratic*. First we record some simple facts about dihedral groups.

Lemma 4.3.13 *Let $C \leq D_{2n}$ be the cyclic subgroup of index two, and say $y \notin C$. Then $o(y) = 2$ and for any $c \in C$, $ycy^{-1} = c^{-1}$.*

If $H \leq D_{2n}$, then H is either cyclic or dihedral. If H is cyclic then either $H \leq C$ or $H \cong \mathbb{Z}/2$, and if H is dihedral then $H = \langle H \cap C, t \rangle$ for any $t \in H - H \cap C$.

The proof is not difficult, and is left to the reader.

Claim 4.3.14 *The number of distinct quadratic subgroups in M is*

$$\frac{o(M)}{12} = \frac{p(p^2 - 1)}{24}.$$

If $o(M)$ is divisible by eight, these subgroups fall into two conjugacy classes of equal size and otherwise they form a single conjugacy class.

Proof. Let $H \leq M$ be a maximal dihedral subgroup, and assume that $4|o(H)$ (so $H \cong D_{2r}$ if r is even and $H \cong D_{2q}$ if q is even; recall that $(r, q) = 1$). Let $C \leq H$ be the cyclic subgroup of index two, and let $t \in C$ be the unique element (in C) of order two. Then if $y \in H - C$, Lemma 4.3.13 shows that $\langle y, t \rangle$ is quadratic. Every quadratic subgroup of H may be obtained in this manner, since by Lemma 4.3.13, any quadratic subgroup $Q \leq H$ contains t and also contains two elements of $H - C$. So as y ranges over $H - C$ the subgroups $\langle y, t \rangle$ range over all quadratic subgroups of H , and each quadratic subgroup is counted twice. Thus there are $\frac{o(C)}{2}$ quadratic subgroups in H .

Next we claim that each quadratic subgroup $Q \leq M$ lies in exactly three conjugates of H , i.e. in exactly three copies of D_{2r} or D_{2q} , depending on the parity of r and q . This follows from Theorem 4.3.3: each of the three non-trivial elements in Q lies in a different conjugate of C , and since Q is abelian Q lies in the corresponding conjugates of $H = N_M(C)$. Certainly Q cannot lie in a fourth conjugate of H , since then two of the corresponding conjugates of C would intersect non-trivially.

The total number of quadratic subgroups of M is now seen to be

$$\frac{\frac{o(C)}{2} c_M(H)}{3}$$

where $c_M(H)$ denotes the number of conjugates of H in M . We have either $o(C) = r = \frac{p+1}{2}$ and $c_M(H) = pq = \frac{p(p-1)}{2}$; or $o(C) = q$ and $c_M(H) = pr = \frac{p(p+1)}{2}$. In either case $o(C)c_M(H) = \frac{p(p^2-1)}{4}$ and the total number of quadratic subgroups is

$$\frac{p(p^2 - 1)}{24} = \frac{o(M)}{12}.$$

If $o(M)$ is not divisible by eight, the quadratic subgroups are actually Sylow 2-subgroups of M and hence form a single conjugacy class.

Now assume that $8|o(M)$. Then $8|o(H)$ and we see that any quadratic subgroup $Q \leq H$ is normalized by a copy of D_8 lying in H . We claim that in

fact, $N_H(Q) \cong D_8$. This follows easily from the Lemma 4.3.13 and the fact that D_{16} does not contain a normal copy of D_4 . Note that this implies that $8|o(N_M(Q))$.

The above argument shows that Q has $\frac{o(H)}{8}$ conjugates in H and since there are twice this many quadratic subgroups in H , the quadratic subgroups of H fall into two conjugacy classes of equal size. Since each quadratic subgroup is contained in a conjugate of H , every quadratic subgroup in M is conjugate to a quadratic subgroup in H . This means that there are at most two conjugacy classes of quadratic subgroups in M .

If all the quadratic subgroups in M were conjugate, then by Lemma 4.3.4 the normalizer of any quadratic subgroup would have order 12, contradicting the fact (mentioned above) that $8|o(N_M(Q))$. Thus there are indeed two conjugacy classes of quadratic subgroups, as desired. $\square$

Our next goal will be to determine the normalizer of a quadratic subgroup of M .

Theorem 4.3.15 *Let $Q \leq M$ be a quadratic subgroup. If $o(M)$ is divisible by eight then $N_Q(M) \cong S_4$ and otherwise $N_Q(M) \cong A_4$.*

Proof. Let $N = N_Q(M)$ and let H be a maximal dihedral subgroup of M containing Q . First we compute $o(N)$. If $o(M)$ is not divisible by eight, Claim 4.3.14 and Lemma 4.3.4 show that $o(N) = 12$, and in the same manner we find $o(N) = 24$ if eight divides $o(M)$.

Now, in the case where $o(N) = 12$, consider an element $s \in N$ of order three. Let $C \leq H$ be the cyclic subgroup of index two. Then by Lemma 4.3.13, $Q \cap C \cong \mathbb{Z}/2$. Let $a \in Q \cap C$ be the non-trivial element. If s commutes with a , then $s \in N(\langle a \rangle) = H$, and if s commuted with another non-trivial element $b \in Q$, s would lie in the intersection of H and one of its conjugates. This is impossible (by Theorem 4.3.3), since then s would lie in C and one of its conjugates. Thus N is a non-abelian group of order twelve containing a normal quadratic subgroup, and any such group is isomorphic to A_4 (Dummit and Foote [9, p. 170]).

Next we must show that when $8|o(M)$, $N \cong S_4$. Since $o(N) = 24$ in this case, we again have an element $s \in N$ of order three, and the same argument as above shows that $\langle s, Q \rangle \cong A_4$. Now, A_4 has four subgroups of order three, and thus there are at least four subgroups of order three in N . In fact, since these subgroups are Sylow 3-subgroups, there can be no more than four of

them (the number of Sylow 3-subgroups is equivalent to 1 (mod 3) and must divide 24).

Consider the action of N (by conjugation) on its four subgroups of order three. This action induces a homomorphism $f : N \rightarrow S_4$, and we claim that this map is an injection. Since $o(N) = o(S_4)$, this will complete the proof.

Let $T = \ker(f)$. Then T is the intersection of the normalizers of the Sylow 3-subgroups of N , so T is characteristic (and hence normal) in N . No (non-trivial) element of $\langle s, Q \rangle \cong A_4$ normalizes each of these subgroups, so $T \cap \langle s, Q \rangle = \{1\}$ and hence

$$\langle s, T, Q \rangle \cong T \times \langle s, Q \rangle \cong T \times A_4 \leq N.$$

Since $o(N) = 24$, we must have $o(T) \leq 2$. If $o(T) = 2$, then $\langle Q, T \rangle \cong D_4 \times \mathbb{Z}/2$. This group is neither cyclic nor dihedral, contradicting the fact that any 2-group in M lies in a copy of D_{2r} or D_{2q} and hence is either cyclic or dihedral. This contradiction shows that f is injective, completing the proof. $\square$

There is one more type of subgroup which sometimes appears in M .

Theorem 4.3.16 *If $p \equiv \pm 1 \pmod{5}$ then M contains exactly $\frac{o(M)}{30}$ subgroups isomorphic to A_5 , falling into two conjugacy classes of equal size.*

Since we are mainly interested in the case $p = 7$, we refer to Burnside [8, §324] for the proof.

The subgroups determined thus far exhaust all the subgroups of M . We will prove this fact in two steps.

Theorem 4.3.17 *Let $H \leq M$ be a subgroup whose order is divisible by p . Then either $H = M$ or $H \leq \text{Stab}(x)$ for some $x \in \mathbb{F}_p \cup \{\infty\}$, i.e. H lies in the normalizer of a Sylow p -subgroup.*

Proof. Since $o(H)$ is divisible by p , there must be a subgroup $P \leq H$ of order p . This subgroup is in fact a Sylow p -subgroup of H , since p^2 does not divide the order of M . The number of Sylow p -subgroups of H is equivalent to 1 (mod p), and hence must be either 1 or $p + 1$ (since M contains just $p + 1$ subgroups of order p). Note that the set of Sylow p -subgroups of M generates a normal subgroup, and since M is simple they must generate the whole group. Thus either H contains a unique (normal) subgroup of order p or $H = M$. $\square$

Theorem 4.3.18 *If $H \leq M$ is a subgroup whose order is not divisible by p , then one of the following is true:*

1. H is cyclic;
2. $H \cong D_{2q'}$ for some $q'|q$ and $H \leq \text{Stab}(x, y)$ for some $x, y \in \mathbb{F}_p \cup \{\infty\}$;
3. $H \cong D_{2r'}$ for some $r'|r$ and H is contained in a copy of D_{2r} ;
4. $H \cong A_4$;
5. $H \cong S_4$;
6. $H \cong A_5$.

Note that all possible cyclic subgroups of M were already determined in Theorem 4.3.3. All dihedral subgroups were determined at this point as well, since any dihedral subgroup lies in the normalizer of its cyclic subgroup of index two. Thus all that remains to be proved is that any subgroup of M has one of the isomorphism types listed above. (Note that A_4 is the only proper subgroup of S_4 or A_5 which is neither cyclic nor dihedral).

Proof. Consider a subgroup $H \leq M$ whose order is not divisible by p . Let $n = o(H)$. We will break H down into conjugacy classes of cyclic subgroups, beginning with the largest cyclic subgroup in H .

Let h_1 be an element of maximum order in H , and let $q_1 = o(h_1)$. By Theorem 4.3.3, $N_H(\langle h_1 \rangle)$ is either cyclic or dihedral, and by maximality of q_1 we see that $o(N_H(\langle h_1 \rangle))$ is either q_1 or $2q_1$. Thus the number of conjugates of $\langle h_1 \rangle$ in H is either $\frac{n}{q_1}$ or $\frac{n}{2q_1}$. By Theorem 4.3.3, no two conjugates of $\langle h_1 \rangle$ may intersect non-trivially. Thus we see that the total number non-trivial elements of H lying in conjugates of $\langle h_1 \rangle$ is

$$\frac{n(q_1 - 1)}{\epsilon_1 q_1},$$

where $\epsilon_1 \in \{1, 2\}$.

Consider the set

$$S_1 = \{h \in H : h \notin \langle h_1 \rangle^z \text{ for any } z \in M\}$$

and let $h_2 \in S_1$ be an element with maximum order q_2 . As before, the total number of non-trivial elements of H lying in conjugates of $\langle h_2 \rangle$ is

$$\frac{n(q_2 - 1)}{\epsilon_2 q_2},$$

where $\epsilon_2 \in \{1, 2\}$. We claim that in fact, none of these elements lie in conjugates of $\langle h_1 \rangle$. Theorem 4.3.3 shows that if two cyclic subgroups of M intersect non-trivially, one must be contained in the other. Thus if $\langle h_2 \rangle^y \cap \langle h_1 \rangle^z$ were non-trivial for some $y, z \in M$, then we would have $\langle h_2 \rangle^y \leq \langle h_1 \rangle^z$ and $\langle h_2 \rangle \leq \langle h_1 \rangle^{zy^{-1}}$, contradicting the fact that $h_2 \in S_1$.

We may continue in this manner, defining S_2 , h_3 and ϵ_3 in the obvious ways, and eventually we find $S_m = \emptyset$ for some m . At this point we have

$$n = 1 + \sum_{i=1}^m \frac{n(q_i - 1)}{\epsilon_i q_i}, \quad (4.1)$$

or

$$\frac{1}{n} = 1 - \sum_{i=1}^m \frac{q_i - 1}{\epsilon_i q_i}. \quad (4.2)$$

Our goal will be to find all the solutions to this equation, and we will break down our analysis by considering the possible values of the ϵ_i .

Letting $A = \{i : \epsilon_i = 1\}$ and $B = \{i : \epsilon_i = 2\}$, we claim that $|A| \leq 1$. Indeed, noting that $q_i \geq 2$ for each i , Equation 4.2 becomes

$$\begin{aligned} \frac{1}{n} &= 1 - \sum_{i \in A} \frac{q_i - 1}{q_i} - \sum_{j \in B} \frac{q_j - 1}{2q_j} \\ &= 1 - \sum_{i \in A} \left(1 - \frac{1}{q_i}\right) - \sum_{j \in B} \left(\frac{1}{2} - \frac{1}{2q_j}\right) \\ &= 1 - |A| + \sum_{i \in A} \frac{1}{q_i} - \frac{1}{2}|B| + \sum_{j \in B} \frac{1}{2q_j} \\ &\leq 1 - |A| + \frac{1}{2}|A| - \frac{1}{2}|B| + \frac{1}{4}|B| \\ &= 1 - \frac{1}{2}|A| - \frac{1}{4}|B|. \end{aligned}$$

Thus $|A| \leq 1$ since $\frac{1}{n} > 0$, and if $|A| = 1$, then $|B| \leq 1$.

We want to show that $m = |A| + |B| \leq 3$. This will imply that there are only several possible cases to consider. If $m > 2$ then (by the above discussion) we must have $|A| = 0$. If $|A| = 0$, then $\epsilon_i = 2$ for all i , and assuming $m > 3$ gives

$$\begin{aligned}
\frac{1}{n} &= 1 - \frac{q_1 - 1}{2q_1} - \frac{q_2 - 1}{2q_2} - \dots \\
&\leq 1 - \frac{q_1 - 1}{2q_1} - \frac{q_2 - 1}{2q_2} - \frac{q_3 - 1}{2q_3} - \frac{q_4 - 1}{2q_4} \\
&= 1 - \frac{(q_1 - 1)q_2q_3q_4 + (q_2 - 1)q_1q_3q_4 + (q_3 - 1)q_1q_2q_4 + (q_4 - 1)q_1q_2q_3}{2q_1q_2q_3q_4} \\
&= 1 - \frac{q_1q_2q_3 + q_1q_2q_4 + q_1q_3q_4 + q_2q_3q_4 - 4q_1q_2q_3q_4}{2q_1q_2q_3q_4} \\
&= \frac{1}{2q_4} + \frac{1}{2q_3} + \frac{1}{2q_2} + \frac{1}{2q_1} - 1 \\
&\leq \frac{1}{4} + \frac{1}{4} + \frac{1}{4} + \frac{1}{4} - 1 = 0,
\end{aligned}$$

a contradiction.

The various remaining cases can all be dealt with in a manner similar to this computation. In each case we either reach a contradiction or show that any group satisfying the formula must have one of the allowed types. To illustrate the methods, we work two of the cases in full. For the other cases, we refer to Burnside [8, §326].

First, if $m = 1$ and $\epsilon_1 = 1$, Equation 4.1 gives

$$n = 1 + \frac{n(q_1 - 1)}{q_1}$$

and solving for n we find that $n = q_1$. Thus in this case $H = \langle h_1 \rangle$, i.e. H is cyclic of order q_1 .

The second case we will consider is $m = 1$, $\epsilon_1 = 1$, and $\epsilon_2 = 2$. In this case we have

$$\begin{aligned}
\frac{1}{n} &= 1 - \frac{q_1 - 1}{q_1} - \frac{q_2 - 1}{2q_2} \\
&= \frac{2q_1q_2 - 2q_2(q_1 - 1) - q_1(q_2 - 1)}{2q_1q_2}
\end{aligned}$$

$$= \frac{2q_2 + q_1 - q_1q_2}{2q_1q_2}.$$

If $q_2 \geq 3$ then

$$\begin{aligned} \frac{1}{n} &\leq \frac{2q_2 + q_1 - 3q_1}{6q_1} \\ &= \frac{1}{6} - \frac{1}{2} + \frac{q_2}{3q_1} \\ &= \frac{1}{3} \left(\frac{q_2}{q_1} - 1 \right). \end{aligned}$$

But $q_2 \leq q_1$ by construction, so $\frac{q_2}{q_1} - 1 \leq 0$, a contradiction.

So $q_2 = 2$ and we have

$$\begin{aligned} \frac{1}{n} &= 1 - \frac{q_1 - 1}{q_1} - \frac{2 - 1}{4} = 1 - \frac{q_1 - 1}{q_1} - \frac{1}{4} \\ &= \frac{4q_1 - 4(q_1 - 1) - q_1}{4q_1} = \frac{4 - q_1}{4q_1}. \end{aligned}$$

Thus $q_1 < 4$. If $q_1 = 2$, then $\frac{1}{n} = \frac{4-2}{8} = \frac{1}{4}$ and $n = 4$. It is easy to see that no group of order four could have these values for m , ϵ_1 and ϵ_2 . So $q_1 = 3$, and we find $n = 12$. Thus H is a group of order twelve and $h_1 \in H$ is an element of order three with $N_H(\langle h_1 \rangle) = \langle h_1 \rangle$. This means $\langle h_1 \rangle$ has four conjugates in H , and A_4 is the only group of order twelve containing four subgroups of order three (Dummit and Foote [9, p. 170]). $\square$

We now record a corollary which will help us determine the Möbius function of $PSL_2(\mathbb{F}_7)$.

Corollary 4.3.19 *If the order of M is not divisible by 60, then no two subgroups of M intersect in a copy of A_4 .*

Proof. Since $60 = o(A_5)$ does not divide $o(M)$, M does not contain subgroups isomorphic to A_5 . Thus we must simply show that no two subgroups H and K isomorphic to S_4 can intersect in a copy of A_4 . But if $H \cap K \cong A_4$, then $H \cap K$ is normal in both H and K . This is impossible, since H and K are maximal and M is a simple group. $\square$

4.3.2 The Möbius Function of $PSL_2(\mathbb{F}_7)$

We will now describe the theory of Möbius Inversion for finite groups, following Hall's original paper [11]. Most of the proofs are omitted, and the interested reader may refer to Hall's paper. Once the necessary facts have been stated, we will proceed to calculate the Möbius function of $PSL_2(\mathbb{F}_7)$.

Definition 4.3.20 *Let G be a finite group and let $\hat{S}(G)$ denote the poset of all subgroups of G , ordered by inclusion (so $\{1\}, G \in \hat{S}(G)$.) Then the Möbius function of G , $\mu_G : \hat{S}(G) \rightarrow \mathbb{Z}$, is given by*

$$\mu_G(H) = \begin{cases} \tilde{\chi}(\Delta(S(G)_{>H})), & H < G \\ 1, & H = G, \end{cases}$$

where $\tilde{\chi}$ denotes the reduced Euler characteristic and $S(G)_{>1} = S(G)$.

The importance of the Möbius function comes from the following theorem.

Theorem 4.3.21 (Möbius Inversion) *Let $f : \hat{S}(G) \rightarrow \mathbb{Z}$ be any function and let $g : \hat{S}(G) \rightarrow \mathbb{Z}$ be the function*

$$g(H) = \sum_{K \leq H} f(K).$$

Then

$$f(G) = \sum_{H \in \hat{S}(G)} \mu_G(H) g(H).$$

We will use Möbius inversion to count automorphism classes of generating pairs in $PSL_2(\mathbb{F}_7)$. We now take a moment to explain how this works.

An automorphism class of generating pairs is a set of the form

$$\{(\gamma(\alpha), \gamma(\beta)) : \alpha, \beta \in G, \gamma \in \text{Aut}(G)\}.$$

Note that in any automorphism class of generating pairs the orders of the first and second elements remain constant.

Definition 4.3.22 *Let G be a finite group. For $a, b > 1$, let $\Phi_{a,b}(G)$ be the number of automorphism classes of generating pairs (α, β) with $o(\alpha) = a$, $o(\beta) = b$, and let $\phi_{a,b}(G)$ be the number of ordered pairs (α, β) ($\alpha, \beta \in G$) with $o(\alpha) = a$, $o(\beta) = b$ and $\langle \alpha, \beta \rangle = G$.*

Also, let $\sigma_{a,b}(G) = |\{\alpha \in G : o(\alpha) = a\}| |\{\beta \in G : o(\beta) = b\}|$.

Note that for any $a, b > 1$ we have

$$\Phi_{a,b}(G) = \frac{\phi_{a,b}(G)}{o(\text{Aut}(G))}, \quad (4.3)$$

since an automorphism which fixes a set of generators is the identity. We now have the following important application of Theorem 4.3.21.

Corollary 4.3.23 *For any finite group G and any $a, b > 1$,*

$$\Phi_{a,b}(G) = \frac{1}{o(\text{Aut}(G))} \sum_{H \leq G} \mu_G(H) \sigma_{a,b}(H).$$

Proof. This follows immediately from Theorem 4.3.21 and Equation 4.3 after noting that

$$\sum_{H \leq G} \phi_{a,b}(H) = \sigma_{a,b}(G).$$

This latter statement is true because each pair (α, β) generates a unique subgroup of G . $\square$

Before calculating the Möbius function of $PSL_2(\mathbb{F}_7)$, we note some useful facts from Hall [11].

Lemma 4.3.24 *For any $H < G$,*

$$\mu_G(H) = - \sum_{K > H} \mu_G(K),$$

and (equivalently)

$$\sum_{K \geq H} \mu_G(K) = 0.$$

If H is a maximal subgroup of G , then $\mu_G(H) = -1$.

If $H < G$ is not the intersection of a set of maximal subgroups of G , then $\mu_G(H) = 0$.

We now turn to the case $G = PSL_2(\mathbb{F}_7)$. Before beginning the computation of μ_G , we note that any maximal subgroup of G is isomorphic to either S_4 or $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$. This is because in this case, $p = 7, q = 3$ and $r = 4$, so $D_{2q} = D_6 \cong S_3$ and $D_{2r} = D_8$; hence these subgroups sit inside copies of S_4

(following Burnside [8], we will call subgroups isomorphic to S_4 *octahedral*). By Lemma 4.3.24, we need only consider subgroups formed by intersecting copies of $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$ and of S_4 .

Note that the only proper, non-trivial subgroups of $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$ are the normal copy of $\mathbb{Z}/7$ and seven copies of $\mathbb{Z}/3$. Every other proper subgroup of G is a subgroup of S_4 , and each of these is isomorphic to one of the following groups: S_4 , A_4 , D_8 , $D_6 \cong S_3$, D_4 , $\mathbb{Z}/4$, $\mathbb{Z}/3$, $\mathbb{Z}/2$ or the trivial group.

The following observation will be useful: if $H, K \leq G$ and $\exists \gamma \in \text{Aut}(G)$ such that $\gamma(H) = K$, then $\mu_G(H) = \mu_G(K)$ (here G is any finite group). This follows from the fact that γ induces a simplicial isomorphism between $\Delta(S(G)_{>H})$ and $\Delta(S(G)_{>K})$.

Theorem 4.3.3 shows that if $H, K \leq G$ are isomorphic cyclic subgroups, then H and K are conjugate. The same is true if $H \cong K \cong \mathbb{Z}/7 \rtimes \mathbb{Z}/3$, D_6 , or D_8 . Unfortunately, Theorem 4.3.14 tell us that the quadratic subgroups of G fall into two conjugacy classes, and hence the same is true for octahedral subgroups and the subgroups isomorphic to A_4 . Luckily, there is an outer automorphism of G interchanging these conjugacy classes. In order to explain this automorphism, we need the following facts.

Fact 4.3.25 $GL_3(\mathbb{F}_2) \cong PSL_2(\mathbb{F}_7)$.

Proof. The order of $GL_3(\mathbb{F}_2)$ can be calculated in a similar manner to that of $GL_2(\mathbb{F}_p)$ (as in the proof of Claim 4.3.2) and we find that $o(GL_3(\mathbb{F}_2)) = 168$. In addition, note that for any n , $GL_n(\mathbb{F}_2) = SL_n(\mathbb{F}_2)$ since $\mathbb{F}_2$ has only one non-zero element. Also, it is not hard to check that the center of $GL_n(\mathbb{F}_2)$ is trivial, so $GL_3(\mathbb{F}_2) \cong PSL_3(\mathbb{F}_2)$, and hence this group is simple (by Theorem 4.3.1). Up to isomorphism, there is only one simple group of order 168, so we must have $GL_3(\mathbb{F}_2) \cong PSL_2(\mathbb{F}_7)$. (For a proof of this last fact, see Dummit and Foote [9, Exercise 27, p. 215].) $\square$

Fact 4.3.26 For any ring R and any $n > 1$, the map $A \mapsto (A^T)^{-1}$ is an automorphism of $GL_n(R)$. (This map is called the *transpose-inverse*.)

Proof. First, an easy induction shows that $\det(A) = \det(A^T)$, and since $\det(A^{-1}) = -\det(A)$, we see that the transpose-inverse is indeed a *function* from $GL_n(R)$ to $GL_n(R)$. Next, note that if $(A^T)^{-1} = I$, then $(A^T) = I$ and hence $A = I$. So all that remains to be shown is that this map is a homomorphism.

This follows easily from the fact that $(AB)^T = B^T A^T$, as we now have $((AB)^T)^{-1} = (B^T A^T)^{-1} = (A^T)^{-1} (B^T)^{-1}$. $\square$

The group $GL_3(\mathbb{F}_2)$ has a natural action on $\mathbb{F}_2^3 - \{0\}$ via multiplication. We take a moment to study this action.

Lemma 4.3.27 *The action of $GL_3(\mathbb{F}_2)$ on $\mathbb{F}_2^3 - \{0\}$ is 2-transitive, and the stabilizer of any point is isomorphic to S_4 .*

Proof. That this action is 2-transitive depends heavily on the fact that we are working over $\mathbb{F}_2$, and hence the span of any vector $v \in \mathbb{F}_2^3 - \{0\}$ is just $\{0, v\}$. Now, thinking of $GL_3(\mathbb{F}_2)$ as the group of vector-space automorphisms of $\mathbb{F}_2^3$, for any non-zero vectors $v \neq w$ there is a third vector $u \in \mathbb{F}_2^3$ s.t. $\{u, v, w\}$ forms a basis for this vector space. Now we can find an automorphism of $\mathbb{F}_2^3$ sending v to v' and w to w' for any $v' \neq w' \in \mathbb{F}_2^3 - \{0\}$. This shows that the action is 2-transitive, as desired.

To compute the stabilizer of a point, we will think of $GL_3(\mathbb{F}_2)$ as a group of matrices acting by multiplication. Then $\text{Stab}(1, 0, 0)$ is the set of all matrices of the form

$$\begin{bmatrix} 1 & a & b \\ 0 & c & d \\ 0 & e & f \end{bmatrix},$$

and it is easy to see that there are 24 such matrices in $GL_3(\mathbb{F}_2)$. All subgroups of order 24 in $GL_3(\mathbb{F}_2) \cong PSL_2(\mathbb{F}_7)$ are octahedral, so we have $\text{Stab}(1, 0, 0) \cong S_4$ and now the fact that this action is transitive completes the proof. $\square$

Fact 4.3.28 *There is an automorphism of $PSL_2(\mathbb{F}_7)$ which interchanges the two conjugacy classes of octahedral subgroups. This automorphism is induced by the transpose-inverse automorphism of $GL_3(\mathbb{F}_2)$.*

Proof. Lemma 4.3.27 shows that one conjugacy class of octahedral subgroups arises as stabilizers of points in $\mathbb{F}_2^3 - \{0\}$. The second conjugacy class is obtained by applying the transpose-inverse. If $A \in \text{Stab}(1, 0, 0)$, then any easy computation shows that $(A^T)^{-1}$ has the form

$$\begin{bmatrix} 1 & 0 & 0 \\ a & b & c \\ d & e & f \end{bmatrix},$$

for some $a, b, c, d, e, f \in \mathbb{F}_2$. The matrices of this form then form an octahedral subgroup H (note that there are 24 such matrices and transpose-inverse is an automorphism) and it remains to show that this subgroup is not the stabilizer of any point in $\mathbb{F}_2^3$. This is easily shown by direct computation. $\square$

Note that since each quadratic subgroup has an octahedral normalizer, the automorphism described above also interchanges the conjugacy classes of quadratic subgroups. The same is true for the subgroups isomorphic to A_4 .

Theorem 4.3.29 *Let $G = PSL_2(\mathbb{F}_7)$. If $H < G$ then we have:*

$$\mu_G(H) = \begin{cases} \mu_G(H) = -1, & H \cong S_4 \text{ or } \mathbb{Z}/7 \rtimes \mathbb{Z}/3 \\ \mu_G(H) = 1, & H \cong D_8 \text{ or } D_6 \\ \mu_G(H) = 2, & H \cong \mathbb{Z}/3 \\ \mu_G(H) = -4, & H \cong \mathbb{Z}/2 \\ \mu_G(H) = 0, & \text{otherwise.} \end{cases}$$

Proof. If $H \cong S_4$ or $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$, then $\mu_G(H) = -1$ by Lemma 4.3.24.

If $H \cong \mathbb{Z}_7$ then $\mu_G(H) = 0$ because H is not the intersection of two maximal subgroups of G (copies of $\mathbb{Z}/p \rtimes \mathbb{Z}/q$ are stabilizers of points in $\mathbb{F}_p \cup \{\infty\}$, and intersect in copies of $\mathbb{Z}/q$).

If $H \cong \mathbb{Z}_3$, then $H = \text{Stab}(x) \cap \text{Stab}(y)$ for some $x, y \in \mathbb{F}_p \cup \{\infty\}$, and hence $H \leq \text{Stab}(x), \text{Stab}(y)$. Any other subgroups containing a subgroup of order three is isomorphic to D_6 , A_4 or S_4 , and thus must contain $N_G(H) \cong D_6$. By Lemma 4.3.24 we have

$$\mu_G(H) = - \left(\mu_G(\text{Stab}(x)) + \mu_G(\text{Stab}(y)) + \sum_{K \geq N_G(H)} \mu_G(K) \right),$$

and applying the lemma again yields $\mu_G(H) = -(-1 - 1 + 0) = 2$.

If $H \cong A_4$ then $\mu_G(H) = 0$ because H is not an intersection of maximal subgroups (Corollary 4.3.19).

If $H \cong \mathbb{Z}/4$, then any subgroup containing H also contains $N_G(H) \cong D_8$, and hence

$$\mu_G(H) = - \sum_{K \geq N_G(H)} \mu_G(K) = 0.$$

In the following calculations, we will exploit a useful counting formula. Let C and C' be automorphism classes of subgroups of G . Then each element

$K \in C'$ contains the same number of elements $H \in C$ (call this number α) and each element $H \in C$ is contained in the same number of elements of C' (call this number β). We have the following formula for $|C|$:

$$|C| = \frac{|C'|\alpha}{\beta}. \quad (4.4)$$

If $H \cong D_8$, then any subgroup containing H is octahedral. There are $pq = 21$ copies of D_8 in G , forming a single conjugacy class, and there are $\frac{o(G)}{12} = 14$ copies of S_4 forming a single automorphism class. Thus we have $21 = \frac{14 \cdot 3}{\beta}$, and hence $\beta = 2$. Thus $\mu_G(H) = -(-1 - 1 + \mu_G(G)) = 1$. For $H \cong D_6, D_4$ or $\mathbb{Z}/2$, the computation follows similarly.

Finally, Lemma 4.3.24 can be used to show $\mu_G(\{1\}) = 0$. $\square$

Now that we know the Möbius function of G , we can give an explicit form of Corollary 4.3.23. Clearly $\sigma_{a,b}(H)$ depends only on the isomorphism type of H , and we have seen that the same is true of $\mu_G(H)$. By counting the number of subgroups of each isomorphism type we obtain the following inversion formula (for any $a, b > 1$):

$$\begin{aligned} \phi_{a,b}(G) &= \sigma_{a,b}(G) - 14\sigma_{a,b}(S_4) - 8\sigma_{a,b}(\mathbb{Z}/7 \rtimes \mathbb{Z}/3) + 21\sigma_{a,b}(D_8) \\ &\quad + 28\sigma_{a,b}(D_6) + 56\sigma_{a,b}(\mathbb{Z}/3) - 84\sigma_{a,b}(\mathbb{Z}/2). \end{aligned} \quad (4.5)$$

In order to count automorphism classes, we need to know the order of $\text{Aut}(G)$. This can be calculated using Möbius inversion.

Fact 4.3.30 $o(\text{Aut}(PSL_2(\mathbb{F}_7))) = 336$.

Proof. First, note that the group $GL_2(\mathbb{F}_7)$ acts by conjugation on $SL_2(\mathbb{F}_7)$ and induces automorphisms of G (note that I and $-I$ are fixed by conjugation). Thus we have a map $f : GL_2(\mathbb{F}_7) \rightarrow \text{Aut}(G)$. We claim that the kernel of this map is $\{\lambda I : \lambda \in \mathbb{F}_p^*\}$. Certainly these elements act trivially, and a simple matrix computation shows that any element in $\ker(f)$ has this form. This shows that

$$o(\text{Aut}(G)) \geq \frac{o(GL_2(\mathbb{F}_7))}{o(\ker(f))} = \frac{(7^2 - 1)(7^2 - 7)}{6} = 336.$$

To prove that there are no other elements in $\text{Aut}(G)$, we can Equation 4.5 and the observation (made earlier) that $o(\text{Aut}(G))$ divides $\phi_{a,b}$ for any $a, b > 1$. Now, consider $\phi_{2,3}$. The above formula gives

$$\phi_{2,3}(G) = \sigma_{2,3}(G) - 14\sigma_{2,3}(S_4) + 28\sigma_{2,3}(D_6)$$

$$= 21 \cdot 56 - 14 \cdot 9 \cdot 8 + 28 \cdot 3 \cdot 2 = 336,$$

so $o(\text{Aut}(G)) = 336$ as desired. $\square$

Corollary 4.3.31 *For any $a, b > 1$,*

$$\Phi_{a,b}(G) = \frac{\phi_{a,b}(G)}{336},$$

where $\Phi_{a,b}$ denotes the number of automorphism classes of generating pairs with orders a and b .

4.3.3 Simple Connectivity of $\mathcal{C}(PSL_2(\mathbb{F}_7))$

The proof of simple connectivity will be analagous to the second proof for $\mathcal{C}(A_5)$, and will use 2-transitive actions. In order to minimize the amount of computation in the proof, we will now prove two helpful lemmas.

Lemma 4.3.32 *Let $\alpha = \frac{ax+b}{cx+d}$ be any non-trivial element of $PSL_2(\mathbb{F}_7)$. We define the trace-squared of α to be $tr^2(\alpha) = (a+d)^2$ (note that this is the square of the trace of either representative of α in $SL_2(\mathbb{F}_7)$, and thus is well-defined).*

The order of α is determined as follows:

$$o(\alpha) = \begin{cases} 2, & tr^2(\alpha) = 0 \\ 3, & tr^2(\alpha) = 1 \\ 4, & tr^2(\alpha) = 2 \\ 7, & tr^2(\alpha) = 4 \end{cases}$$

Proof. In order to prove this, it will be useful to define the discriminant of a non-trivial element of $G = PSL_2(\mathbb{F}_7)$. If $\alpha = \frac{ax+b}{cx+d} \in G$, we define $disc(\alpha)$ to be the discriminant of the quadratic polynomial $(cx+d)x - (ax+b) = cx^2 + (d-a)x - b$ determined by the equation $\frac{ax+b}{cx+d} = x$, so that $disc(\alpha) = (d-a)^2 - 4(-b)(c) = tr^2(\alpha) - 4$.

We will consider only the elements in $G\text{-Stab}(\infty)$. It is easy to check the result on remaining elements of G . The elements of order seven in $G\text{-Stab}(\infty)$ are exactly those with one fixed point in $\mathbb{F}_7$, i.e. those with $disc(\alpha) = 0$ and $tr^2(\alpha) = 4$. Similarly, elements of order three in $G\text{-Stab}(\infty)$ are those with two fixed points in $\mathbb{F}_7$, i.e. those with $disc(\alpha) = z^2$ for

some $z \in \mathbb{F}_7^*$. Thus $o(\alpha) = 3 \iff \text{disc}(\alpha) \in \{1, 2, 4\} \iff \text{tr}^2(\alpha) \in \{5, 6, 1\} \iff \text{tr}^2(\alpha) = 1$.

Next, let $A \in SL_2(\mathbb{F}_7)$ be a matrix representing $\alpha \in G$, and let $\lambda_1, \lambda_2 \in \bar{\mathbb{F}}_7$ be the eigenvalues of A (where $\bar{\mathbb{F}}_7$ denotes the algebraic closure). Then $o(\alpha) = 2 \implies \lambda_1^2 = \lambda_2^2 = \pm 1$. If $\lambda_1 = \lambda_2$ then $A = \lambda_1 I$ and $\alpha = 1$, so $o(\alpha) = 2 \iff \lambda_1 = -\lambda_2 \iff \text{tr}(A) = \lambda_1 + \lambda_2 = 0 \iff \text{tr}^2(\alpha) = 0$.

Finally, a similar but more lengthy calculation shows that $o(\alpha) = 4 \iff \text{tr}^2(\alpha) = 2$. $\square$

Lemma 4.3.33 *Let $g, h \in PSL_2(\mathbb{F}_7)$ be elements of orders two and three, respectively. Then $\langle g, h \rangle = PSL_2(\mathbb{F}_7) \iff o(gh) = 7$.*

Proof. Certainly, if $o(g) = 2$, $o(h) = 3$ and $o(gh) = 7$, $\langle g, h \rangle = PSL_2(\mathbb{F}_7)$, since no subgroup contains elements of orders two and seven.

In the other direction, recall that there is a unique automorphism class of generators with orders two and three (see the proof of Fact 4.3.30), and thus the result will follow if we show that there exist elements $g, h \in PSL_2(\mathbb{F}_7)$ with $o(g) = 2$, $o(h) = 3$ and $o(gh) = 7$. The elements $g = \frac{x-2}{x-1}$ and $h = \frac{4x}{2}$ satisfy this property, since $\text{tr}^2(g) = 0$, $\text{tr}^2(h) = 1$ (note that $\det(h) = \det(g) = 1$) and $hg = \frac{4\frac{x-2}{x-1}}{2} = \frac{4x-1}{2x-2}$ so $\text{tr}^2(hg) = 4$ and $o(hg) = 7$. $\square$

Theorem 4.3.34 *The coset poset of $PSL_2(\mathbb{F}_7)$ is simply connected.*

Proof. We will show that the minimal cover of $G = PSL_2(\mathbb{F}_7)$ is simply connected. We will employ Theorem 2.1.9, using for a maximal trees the set of all edges $\{1, h\} \in \mathcal{M}(G)$ ($h \in G$). First we will show that all generators of $\pi_1(\mathcal{M}(G))$ corresponding to edges $\{g, h\}$ with $o(g) = 2$ are trivial, and then we will finish the proof by considering 2-transitive actions (note that, since G is non-cyclic, there is an edge between any two vertices in $\mathcal{M}(G)$). As in Theorem 2.1.9, we will denote the generator corresponding to the edge $\{g, h\}$ by (g, h) . [Recall that in the statement of the theorem, there was a generator for each *ordered* edge, but since we have the relation $(u, v) = (v, u)^{-1}$ for each edge $\{u, v\}$, it suffices to show that in each case either (u, v) or (v, u) is trivial.]

We now analyze the various edges $\{g, h\}$ with $o(g) = 2$ appearing in $\mathcal{M}(G)$. First, note that if $\langle g, h \rangle \neq PSL_2(\mathbb{F}_7)$, then the edge (g, h) is trivial because the set $\{1, g, h\}$ is contained in a subgroup. Also, since G is simple two elements of order two cannot generate G and thus if $o(g) = o(h) = 2$,

(g, h) is trivial. We now examine the various possibilities for $o(h)$, and in each case we count the number of automorphism classes of generators (using Equation 4.5) and examine an edge from each class.

$o(g) = 2, o(h) = 3$: Letting $h = \frac{4x}{2}$ and $g = \frac{x-2}{x-1}$, the proof of Lemma 4.3.33 shows that (g, h) is a representative for the unique automorphism class of generators with these orders.

Now, notice that $g(-1) = h(-1) = -2$, so $g \equiv h \pmod{\text{Stab}(-1)}$. Consider the element $z = \frac{3x-2}{-2x-3}$. This element has determinant one and order two, and we have $z(-1) = -2$. Thus $\{g, h, z\}$ is a 2-simplex in $\mathcal{M}(G)$, and we have the relation $(g, h) = (h, z)(g, z)$. Note that $(g, z) = 1$ because these two elements of order two cannot generate G . Thus to prove that $(g, h) = 1$, it suffices to show that $(h, z) = 1$. We have $hz = \frac{4 \cdot \frac{3x-2}{-2x-3}}{2} = \frac{5x-1}{3x+1}$, and hence $o(hz) = 3$. Lemma 4.3.33 now implies that $\langle z, h \rangle \neq G$.

$o(g) = 2, o(h) = 4$: Again, Möbius inversion shows that $\Phi_{2,4}(G) = 1$:

$$\begin{aligned}\phi_{2,4}(G) &= \sigma_{2,4}(G) - 14\sigma_{2,4}(S_4) + 21\sigma_{2,4}(D_8) \\ &= 21 \cdot 42 - 14 \cdot 9 \cdot 6 + 21 \cdot 5 \cdot 2 = 336,\end{aligned}$$

and hence $\Phi_{2,4} = 1$. We claim that the pair $g = \frac{-1}{x}$, $h = \frac{4x+1}{-x}$ is a representative for the unique automorphism class of generators. First, note that these elements each have determinant one and have the correct orders. To show that $\langle g, h \rangle = G$, note that $gh = \frac{-1}{\frac{4x+1}{-x}} = \frac{x}{4x+1}$, so $o(gh) = 7$. Since no proper subgroup of G contains elements of orders two and seven, $\langle g, h \rangle = G$.

Next, note that $g(0) = h(0) = \infty$, and set $z = \frac{-2}{4x}$. Then $\det(z) = 1$, $o(z) = 2$, and $z(0) = \infty$. As in the previous case, if $\langle z, h \rangle \neq G$, then $\{g, h\} = 0$. The proof that $\langle z, h \rangle \neq G$ is also analagous to the previous case: $zh = \frac{-2}{4 \cdot \frac{4x+1}{-x}} = \frac{2x}{2x+4}$ and $o(zh) = 3 \neq o(gh)$. Thus the pair (z, h) does not fall into the unique automorphism class of generators.

$o(g) = 2, o(h) = 7$: This time we find that there are three automorphism classes of generating pairs, since any pair (g, h) with $o(g) = 2$ and $o(h) = 7$ necessarily generates. This means $\phi_{2,a}(G) = 21 \cdot 48 = 1008$ and $\Phi_{2,7} = \frac{1008}{336} = 3$.

Let $h = x + 1$, so $o(h) = 7$, and let $g = \frac{b}{cx}$, so that $o(g) = 2$. We have $hg = \frac{b}{cx} + 1 = \frac{cx+b}{cx}$ and thus $\text{tr}^2(hg) = c^2$. This implies that the pairs

$$(g_1 = \frac{-1}{x}, h), \quad (g_2 = \frac{2}{3x}, h), \quad (g_3 = \frac{3}{2x}, h)$$

represent the three generating automorphism classes.

Next, say there exists an element $z \in \text{Stab}(\infty)$ such that $o(z) = 3$ and $\{h, g_i, z\}$ forms a simplex in $\mathcal{M}(G)$. Then we have $(h, z) = 1$ because $\langle h, z \rangle \leq \text{Stab}(\infty)$ and $(g, z) = 1$ because $o(g_i) = 2$ and $o(z) = 3$ and we have shown above that the generators corresponding to such edges are trivial. We will now attempt to find such elements.

Consider the equations $h(x) = g_i(x)$. These correspond to the equations $x+1 = \frac{b}{-b^{-1}x}$, where $b = -1, 2$ or 3 . Equivalently, [note that ∞ can never be a solution] we are looking for solutions to the quadratic polynomial $x^2 + x + b^2 = 0$, and its discriminant is

$$1 - 4b^2 = \begin{cases} 1 - 4(-1^2) = 4, & b = -1 \\ 1 - 4 \cdot 2^2 = -1, & b = 2 \\ 1 - 4 \cdot 3^2 = 0, & b = 3. \end{cases}$$

Thus solutions exist when $b = -1$ or 3 , but not when $b = 2$.

For the pairs (g_1, h) and (g_3, h) we may now quickly finish the proof by noting that for any $x, y \in \mathbb{F}_7$ there is an element $z \in \text{Stab}(\infty)$ with $o(z) = 3$ and $z(x) = y$, as can be shown by direct computation (it suffices to consider elements of the form $\frac{2x+b}{4}$). Taking into account the previous two paragraphs, this shows that $(g_1, h) = (g_3, h) = 1$.

Finally, we must show that $(g_2, h) = (\frac{2}{3x}, x+1) = 1$. Letting $z = \frac{2x}{4}$, we have $o(z) = 3$ and $z \in \text{Stab}(\infty)$, so it suffices to show that these three elements lie in a proper coset, i.e. that $\langle z^{-1}g_2, z^{-1}h \rangle \neq G$. We have $z^{-1} = \frac{4x}{2}$, so $z^{-1}g_2 = \frac{4 \cdot \frac{2}{3x}}{2x} = \frac{1}{-x}$ and hence $o(z^{-1}g_2) = 2$. Next, $z^{-1}h = \frac{4(x+1)}{2}$ so $o(z^{-1}h) = 3$. Also, $z^{-1}g_2z^{-1}h = \frac{1}{-\frac{4x+4}{2}} = \frac{2}{3x-3}$, so $o(z^{-1}g_2z^{-1}h) = 4$ and hence $\langle z^{-1}g_2, z^{-1}h \rangle \neq G$.

We have now shown that all edges $\{g, h\}$ with $o(g) = 2$ correspond to trivial generators. We will complete the proof of simple connectivity by considering 2-transitive actions of G . Note that since G is not cyclic, any two elements of G are equivalent modulo some maximal subgroup. Thus it will suffice to show that there is an element of order two in each coset kM where $k \in G$ and $M \leq G$ is maximal. We will do this by showing that in each case the action of G on the cosets G/M is two-transitive and that some element of order two acts non-trivially.

First, recall that each subgroup isomorphic to $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$ is the stabilizer of a point in $\mathbb{F}_7 \cup \{\infty\}$, and this action is two-transitive (Lemma 4.3.7). Also,

since no element of G other than the identity acts trivially on $\mathbb{F}_7 \cup \{\infty\}$, there are certainly elements of order two which act non-trivially.

Next, Lemma 4.3.27 shows that the action of G on one of its conjugacy classes of octahedral subgroups is 2-transitive, and Fact 4.3.28 shows that there is an automorphism interchanging the two conjugacy classes, so in fact G acts 2-transitively on each conjugacy class of octahedral subgroups. The stabilizer of a subgroup is just its normalizer, and the octahedral subgroups are self-normalizing (because they are maximal and G is simple). Finally, we need to check that there are elements of order two which act non-trivially. This follows immediately from the fact that in $GL_3(\mathbb{F}_2)$, only the identity acts trivially on $\mathbb{F}_2^3 - \{0\}$ (and this action has octahedral stabilizers). $\square$

There are essentially two barriers preventing the extension of this result to $PSL_2(\mathbb{F}_p)$ for primes $p > 7$. First, it is not clear how to generalize the ad hoc portion of the proof, in which we showed that all edges $\{g, h\}$ with $o(g) = 2$ corresponded to trivial generators of $\pi_1(\mathcal{M}(G))$. Also, for large p the only 2-transitive action of $PSL_2(\mathbb{F}_p)$ is its standard action on $\mathbb{F}_p \cup \{\infty\}$. This follows from [15, Exercises 38 and 39, p. 58], and the essential reason is that the subgroups of $PSL_2(\mathbb{F}_p)$ are too small, and thus have large index. (Whenever a group G acts 2-transitively on a set X , we get a transitive action of G on the set $X \times X - \{(x, x) : x \in X\}$, and thus this set can be no larger than G . It is possible that there are other small primes for which one gets multiple 2-transitive actions.)

There are a few more facts about the homotopy-type of $\mathcal{C}(PSL_2(\mathbb{F}_7))$ that may be proven by elementary methods. In particular, it is possible to show that $\mathcal{C}(PSL_2(\mathbb{F}_7))$ is homotopy equivalent to a three-dimensional complex. We will briefly explain the process.

First, note that any chain of length five lies under an octahedral subgroup, since the only other type of maximal subgroup is $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$. It is easy to check that every chain of length five contains a coset xH where $H \cong D_4$ or $H \cong \mathbb{Z}/4$. But cosets of copies of D_4 and $\mathbb{Z}/4$ may be removed from $\mathcal{C}(PSL_2(\mathbb{F}_7))$ without changing the homotopy type, as can be shown by applying Quillen's Theorem to the inclusion map (see the proof of Claim 4.2.4). This shows that $\mathcal{C}(PSL_2(\mathbb{F}_7))$ is homotopy equivalent to a three-dimensional complex.

We can now show that $H_2(\mathcal{C}(PSL_2(\mathbb{F}_7)))$ is non-trivial, and in fact has rank at least $17 \cdot 168$. This follows immediately from a computation of the Euler characteristic of $\mathcal{C}(PSL_2(\mathbb{F}_7))$. It is shown in Brown [7, Table I] that

$\tilde{\chi}(\mathcal{C}(PSL_2(\mathbb{F}_7))) = 17 \cdot 168$, and since the only even dimension in which $\mathcal{C}(PSL_2(\mathbb{F}_7))$ has homology is dimension two, the result follows.

It is possible to eliminate various other simplices from $\Delta(\mathcal{C}(G))$ without changing the homotopy type. For example, all cosets of all subgroups isomorphic to A_4 and $\mathbb{Z}/7$ may be removed (by applying Quillen's Theorem to the inclusion map), and after removing the cosets xA_4 we may remove all simplices of the form $\{x\} \subset xC_3 \subset xD_6 \subset xS_4$ or $\{x\} \subset xC_3 \subset xS_4$ (where C_3 denotes a cyclic subgroup of order three). The latter type of chain is contained in a unique maximal chain (obtained by adding the coset $xN(C_3)$) and thus removing these simplices constitutes an "elementary collapse" (see Brown [6]). Unfortunately, it does not seem possible to eliminate all chains of length four, and thus we are unable to prove that $\mathcal{C}(PSL_2(\mathbb{F}_7))$ is homotopy equivalent to a wedge of two-spheres. In fact, we conjecture that $H_3(\mathcal{C}(PSL_2(\mathbb{F}_7)))$ is non-trivial (see Chapter 5).

Chapter 5

Conjectures and Directions for Further Research

In this final chapter, we briefly discuss some conjectures about the coset poset and some potential directions further research could take. Other such remarks have been scattered through the previous chapters, but these seemed not to fit in elsewhere.

There appear to be certain concrete connections between the subgroup poset and the coset poset of a finite group G ; in some sense, the structure of the coset poset depends heavily on the structure of the subgroup poset.

There is a general theme in the proofs that the coset posets A_5 and $PSL_2(\mathbb{F}_7)$ are homotopy equivalent to lower dimensional subposets: we may remove cosets from $\mathcal{C}(G)$ without changing the homotopy type whenever the corresponding subgroups can be removed from $S(G)$ (without changing the homotopy type). Initially, the coset poset has dimension one more than the dimension of $S(G)$ (due to the fact that singleton cosets are included in $\mathcal{C}(G)$), and as we remove cosets in this fashion, the relationship between dimensions remains intact. This suggests the idea that the coset poset should have “homotopy dimension” exactly one more than the “homotopy dimension” of the subgroup poset, where by homotopy dimension we mean the lowest dimension of a simplicial complex with the same homotopy type. (In some cases, the subgroup poset is contractible and this relationship does not hold.) For solvable groups with non-contractible subgroup posets, this relationship follows immediately from Brown’s determination of the homotopy type of $\mathcal{C}(G)$ (Theorem 3.3.2) and the following similar result of Kratzer and

Thévenaz [14, Corollaire 4.10]:

Theorem 5.0.1 *If G is a finite solvable group, then $S(G)$ is homotopy equivalent to a wedge of spheres of dimension $d - 2$, where d is the length of a chief series for G (in the sense of Theorem 3.3.2). The number of spheres is zero if and only if some subgroup in a chief series for G does not have a complement.*

It is worth mentioning at this point that the coset poset can be viewed as a homotopy colimit, in the sense of [23]. The underlying category is the poset $S(G) \cup \{1\}$, and the space above a subgroup H is a discrete set of $(G : H)$ points (thought of as the cosets of H in G). The map associated to the inclusion $H < K$ is just $xH \mapsto xK$, and the fact that this diagram has homotopy colimit homeomorphic to $\mathcal{C}(G)$ is an immediate consequence of [23, Proposition 4.1]. Originally I had hoped that, given a poset $P \simeq S(G)$, one could find a diagram over $C(P)$ (the cone on P , i.e. P with an extra element 0 satisfying $0 < p$ for each $p \in P$) with homotopy colimit $H \simeq \mathcal{C}(G)$. The desired dimension result would then follow immediately from [23, Proposition 4.1]. Unfortunately, the literature on homotopy colimits does not seem to contain results of this nature.

In Theorem 3.2.3, we saw that in certain cases there is a surjection from $H_{n+1}(\mathcal{C}(G))$ to $H_n(S(G))$, and I wonder whether the hypothesis (that $\mathcal{C}(G)_g$ has trivial n -dimensional homology) is truly necessary. It is almost certain that the hypothesis is not always satisfied (for example I doubt that $PSL_2(\mathbb{F}_7)$ satisfies the hypothesis when $n = 2$), but the theorem may remain true for other reasons. In particular, one ought to be able to settle this question for solvable groups, due to the two theorems discussed above.

When applied to the coset poset of $G = PSL_2(\mathbb{F}_7)$, these conjectures leads me to believe that $H_3(\mathcal{C}(G))$ is probably non-zero, because $H_2(S(G))$ is non-zero. The latter fact follows from two previous observations and a result of Shareshian: from Theorem 4.3.29 we know that the reduced Euler characteristic of $S(G)$ is zero, and Shareshian [21, Lemma 3.11] has shown that $H_1(S(G)) \neq 0$ (this follows easily by considering the graph formed by subgroups of type $\mathbb{Z}/3$ and $\mathbb{Z}/7 \rtimes \mathbb{Z}/3$ and noting that the edges of this graph are maximal simplices in $S(G)$). The proof (sketched at the end of Section 4.3.3) that $\mathcal{C}(G)$ is homotopy equivalent to a three-dimensional complex also shows that $S(PSL_2(\mathbb{F}_7))$ is homotopy equivalent to a two-dimensional

complex, and thus $H_2(S(G))$ must be non-zero in order for $\tilde{\chi}(S(G))$ to be zero.

Even if $H_3(\mathcal{C}(G))$ is zero, our first conjecture about dimensions suggests that $\mathcal{C}(G)$ is not a bouquet of two-spheres, since $S(G)$ is not homotopy equivalent to a one-dimensional complex. This would yield the only known example of a group whose coset poset is not a bouquet of equal-dimensional spheres. (I do not know whether there are other examples of groups G with $S(G)$ not a bouquet of spheres; it is possible that $PSL_2(\mathbb{F}_7)$ is the only such group known.)

On a different note, it would be interesting to study the coset poset of an infinite group. Many of the results in this thesis probably extend (in some manner) to this more general setting. For example, the Nerve Theorem holds for infinite simplicial complexes (under appropriate conditions) and this should allow one to define the minimal cover of an infinite group.

Bibliography

- [1] M. Aschbacher. Simple connectivity of p -group complexes. *Israel J. Math.*, **82**:1–43, 1993.
- [2] M. Aschbacher and S. D. Smith. On Quillen’s conjecture for the p -groups complex. *Ann. of Math. (2)*, **137**:473–529, 1993.
- [3] A. Björner. Topological methods. In *Handbook of Combinatorics, Vol. 2*, pages 1819–1872. Elsevier, Amsterdam, 1995.
- [4] A. Björner and J. W. Walker. A homotopy complementation formula for partially ordered sets. *European J. Combinatorics*, **4**:11–19, 1983.
- [5] K. S. Brown. *Cohomology of Groups*. Springer-Verlag, New York, 1982.
- [6] K. S. Brown. The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem. In *Algorithms and Classification in Combinatorial Group Theory (Berkeley, CA, 1989)*, volume 23 of *Math. Sci. Res. Inst. Publ.*, pages 137–163. Springer, New York, 1992.
- [7] K. S. Brown. The coset poset and probabilistic zeta function of a finite group. *Journal of Algebra*, **225**:989–1012, 2000.
- [8] W. Burnside. *Theory of Groups of Finite Order*. Dover, New York, 2nd edition, 1955.
- [9] D. S. Dummit and R. M. Foote. *Abstract Algebra*. Wiley, New York, 2nd edition, 1999.
- [10] S. Eilenberg and N. Steenrod. *Foundations of Algebraic Topology*. Princeton Univ. Press, Princeton, 1952.

- [11] P. Hall. The Eulerian functions of a group. *Quart. J. Math.*, **7**:134–151, 1936.
- [12] A. Hatcher. *Algebraic Topology*. Cambridge Univ. Press, Cambridge, UK, 2002.
- [13] I. N. Herstein. A remark on finite groups. *Proc. Amer. Math. Soc.*, **9**:255–257, 1958.
- [14] C. Kratzer and J. Thévenaz. Type d’homotopie des treillis et treillis des sous-groupes d’un groupe fini. *Comment. Math. Helv.*, **60**:85–106, 1985.
- [15] S. Lang. *Algebra*. Addison-Wesley, Menlo Park, CA, 2nd edition, 1984.
- [16] J. W. Milnor. Construction of universal bundles. II. *Ann. of Math. (2)*, **63**, 1956.
- [17] J. Pulkus and V. Welker. On the homotopy type of the p -subgroup complex for finite solvable groups. *J. Austral. Math. Soc. Ser. A*, **69**:212–228, 2000.
- [18] D. Quillen. Homotopy properties of the poset of nontrivial p -subgroups of a group. *Advances in Mathematics*, **28**:101–128, 1978.
- [19] D. J. S. Robinson. *A Course in the Theory of Groups*. Springer-Verlag, New York, 1996.
- [20] Y. Segev and P. Webb. Extensions of G -posets and Quillen’s complex. *J. Austral. Math. Soc. Ser. A*, **57**:60–75, 1994.
- [21] J. Shareshian. On the shellability of the order complex of the subgroup lattice of a finite group. *Trans. Amer. Math. Soc.*, **353**:2689–2703, 2001.
- [22] J. Thévenaz. Permutation representations arising from simplicial complexes. *J. Combin. Theory Ser. A*, **46**:121–155, 1987.
- [23] V. Welker, G. Ziegler, and R. T. Živaljević. Homotopy colimits - comparison lemmas for combinatorial applications. *J. reine angew. Math.*, **509**:117–149, 1999.